

- Adams JD, Jr., Klaidman LK (1993) Acrolein-induced oxygen radical formation. *Free Radic. Biol. Med.*, **15**: 187-193.
- Agbaga MP, Brush RS, Mandal MN, Henry K, Elliott MH, Anderson RE (2008) Role of Stargardt-3 macular dystrophy protein (ELOVL4) in the biosynthesis of very long chain fatty acids. *Proc. Natl. Acad. Sci. U. S. A.*, **105**: 12843-12848.
- Alderson NL, Rembiesa BM, Walla MD, Bielawska A, Bielawski J, Hama H (2004) The human FA2H gene encodes a fatty acid 2-hydroxylase. *J. Biol. Chem.*, **279**: 48562-48568.
- Alderson NL, Walla MD, Hama H (2005) A novel method for the measurement of in vitro fatty acid 2-hydroxylase activity by gas chromatography-mass spectrometry. *J. Lipid Res.*, **46**: 1569-1575.
- Anderson RA, Joyce C, Davis M, Reagan JW, Clark M, Shelness GS, Rudel LL (1998) Identification of a form of acyl-CoA:cholesterol acyltransferase specific to liver and intestine in nonhuman primates. *J. Biol. Chem.*, **273**: 26747-26754.
- Andrieu MN, Kurtman C, Hicsonmez A, Ozbilgin K, Eser E, Erdemli E (2005) In vivo study to evaluate the protective effects of amifostine on radiation-induced damage of testis tissue. *Oncology*, **69**: 44-51.
- Arenas MI, Lobo MV, Caso E, Huerta L, Paniagua R, Martin-Hidalgo MA (2004) Normal and pathological human testes express hormone-sensitive lipase and the lipid receptors CLA-1/SR-BI and CD36. *Hum. Pathol.*, **35**: 34-42.
- Aveladano MI (1987) A novel group of very long chain polyenoic fatty acids in dipolyunsaturated phosphatidylcholines from vertebrate retina. *J. Biol. Chem.*, **262**: 1172-1179.
- Aveladano MI, Robinson BS, Johnson DW, Poulos A (1993) Long and very long chain polyunsaturated fatty acids of the n-6 series in rat seminiferous tubules. Active desaturation of 24:4n-6 to 24:5n-6 and concomitant formation of odd and even chain tetraenoic and pentaenoic fatty acids up to C32. *J. Biol. Chem.*, **268**: 11663-11669.
- Aveladano MI, Rotstein NP, Vermouth NT (1992a) Lipid remodelling during epididymal maturation of rat spermatozoa. Enrichment in plasmenylcholines containing long-chain polyenoic fatty acids of the n-9 series. *Biochem. J.*, **283 (Pt 1)**: 235-241.
- Aveladano MI, Rotstein NP, Vermouth NT (1992b) Occurrence of long and very long polyenoic fatty acids of the n-9 series in rat spermatozoa. *Lipids*, **27**: 676-680.
- Aveladano MI, Sprecher H (1987) Very long chain (C24 to C36) polyenoic fatty acids of the n-3 and n-6 series in dipolyunsaturated phosphatidylcholines from bovine retina. *J. Biol. Chem.*, **262**: 1180-1186.
- Ayyagari R, Zhang K, Hutchinson A, Yu Z, Swaroop A, Kakuk LE, Seddon JM, Bernstein PS, Lewis RA, Tammur J, Yang Z, Li Y, Zhang H, Yashar BM, Liu J, Petrukhin K, Sieving PA, Allikmets R (2001) Evaluation of the ELOVL4 gene in patients with age-related macular degeneration. *Ophthalmic Genet.*, **22**: 233-239.
- Barenholz Y (2004) Sphingomyelin and cholesterol: from membrane biophysics and rafts to potential medical applications. *Subcell. Biochem.*, **37**: 167-215.
- Baum JS, St George JP, McCall K (2005) Programmed cell death in the germline. *Semin. Cell Dev. Biol.*, **16**: 245-259.
- Beckman JK, Coniglio JG (1979) A comparative study of the lipid composition of isolated rat Sertoli and germinal cells. *Lipids*, **14**: 262-267.
- Beckman JK, Coniglio JG (1980) The metabolism of polyunsaturated fatty acids in rat Sertoli and germinal cells. *Lipids*, **15**: 389-394.
- Beckman JK, Gray ME, Coniglio JG (1978) The lipid composition of isolated rat spermatids and spermatocytes. *Biochim. Biophys. Acta*, **530**: 367-374.
- Bellve AR, Cavicchia JC, Millette CF, O'Brien DA, Bhatnagar YM, Dym M (1977) Spermatogenic cells of the prepuberal mouse. Isolation and morphological characterization. *J. Cell Biol.*, **74**: 68-85.

- Birbes H, El BS, Hannun YA, Obeid LM (2001) Selective hydrolysis of a mitochondrial pool of sphingomyelin induces apoptosis. *FASEB J.*, **15**: 2669-2679.
- BLIGH EG, DYER WJ (1959) A rapid method of total lipid extraction and purification. *Can. J. Biochem. Physiol.*, **37**: 911-917.
- Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal. Biochem.*, **72**: 248-254.
- Brenner RR (1977) Comparative fatty acid composition of lipids from tubular and interstitial cells of rat testicles. *Acta Physiol Lat. Am.*, **27**: 203-206.
- Brewis IA, Moore HD, Fraser LR, Holt WV, Baldi E, Luconi M, Gadella BM, Ford WC, Harrison RA (2005) Molecular mechanisms during sperm capacitation. *Hum. Fertil. (Camb.)*, **8**: 253-261.
- Bridges RB, Coniglio JG (1970) The metabolism of 4,7,10,13,16-(5-14C)docosapentaenoic acid in the testis of the rat. *Biochim. Biophys. Acta*, **218**: 29-35.
- Brown ER, Subbaiah PV (1994) Differential effects of eicosapentaenoic acid and docosahexaenoic acid on human skin fibroblasts. *Lipids*, **29**: 825-829.
- Butler A, Gordon RE, Gatt S, Schuchman EH (2007) Sperm abnormalities in heterozygous acid sphingomyelinase knockout mice reveal a novel approach for the prevention of genetic diseases. *Am. J. Pathol.*, **170**: 2077-2088.
- Butler A, He X, Gordon RE, Wu HS, Gatt S, Schuchman EH (2002) Reproductive pathology and sperm physiology in acid sphingomyelinase-deficient mice. *Am. J. Pathol.*, **161**: 1061-1075.
- Cai L, Hales BF, Robaire B (1997) Induction of apoptosis in the germ cells of adult male rats after exposure to cyclophosphamide. *Biol. Reprod.*, **56**: 1490-1497.
- Cameron DJ, Tong Z, Yang Z, Kaminoh J, Kamiyah S, Chen H, Zeng J, Chen Y, Luo L, Zhang K (2007) Essential role of Elovl4 in very long chain fatty acid synthesis, skin permeability barrier function, and neonatal survival. *Int. J. Biol. Sci.*, **3**: 111-119.
- Cerec V, Piquet-Pellorce C, Aly HA, Touzalin AM, Jegou B, Bauche F (2007) Multiple pathways for cationic amino acid transport in rat seminiferous tubule cells. *Biol. Reprod.*, **76**: 241-249.
- Chao CC, Ma YS, Stadtman ER (1997) Modification of protein surface hydrophobicity and methionine oxidation by oxidative systems. *Proc. Natl. Acad. Sci. U. S. A.*, **94**: 2969-2974.
- Chemes H (1986) The phagocytic function of Sertoli cells: a morphological, biochemical, and endocrinological study of lysosomes and acid phosphatase localization in the rat testis. *Endocrinology*, **119**: 1673-1681.
- Chistie, WW (1982) Lipid analysis, 2nd edition. *Pergamon Press*
- Codrington AM, Hales BF, Robaire B (2004) Spermiogenic germ cell phase-specific DNA damage following cyclophosphamide exposure. *J. Androl*, **25**: 354-362.
- Coniglio JG (1994) Testicular lipids. *Prog. Lipid Res.*, **33**: 387-401.
- Coniglio JG, Grogan WM, Jr., Rhamy RK (1974) Lipids of human testes removed at orchidectomy. *J. Reprod. Fertil.*, **41**: 67-73.
- Coniglio JG, Harris DG (1977) Fatty acid composition of selected organs of gerbils maintained on a fat-deficient or fat-supplemented diet. *Lipids*, **12**: 741-746.
- Connor WE, Lin DS, Wolf DP, Alexander M (1998) Uneven distribution of desmosterol and docosahexaenoic acid in the heads and tails of monkey sperm. *J. Lipid Res.*, **39**: 1404-1411.
- Contreras FX, Basanez G, Alonso A, Herrmann A, Goni FM (2005) Asymmetric addition of ceramides but not dihydroceramides promotes transbilayer (flip-flop) lipid motion in membranes. *Biophys. J.*, **88**: 348-359.
- Cremesti AE, Goni FM, Kolesnick R (2002) Role of sphingomyelinase and ceramide in modulating rafts: do biophysical properties determine biologic outcome? *FEBS Lett.*, **531**: 47-53.

- Crook TR, Souhami RL, McLean AE (1986) Cytotoxicity, DNA cross-linking, and single strand breaks induced by activated cyclophosphamide and acrolein in human leukemia cells. *Cancer Res.*, **46**: 5029-5034.
- Cross NL (2000) Sphingomyelin modulates capacitation of human sperm in vitro. *Biol. Reprod.*, **63**: 1129-1134.
- Cutler RG, Kelly J, Storie K, Pedersen WA, Tammara A, Hatanpaa K, Troncoso JC, Mattson MP (2004) Involvement of oxidative stress-induced abnormalities in ceramide and cholesterol metabolism in brain aging and Alzheimer's disease. *Proc. Natl. Acad. Sci. U. S. A.*, **101**: 2070-2075.
- Dakhova O, O'Day D, Kinet N, Yucer N, Wiese M, Shetty G, Ducy P (2008) Dkk1 regulates post-pubertal spermatocyte apoptosis and testosterone production. *Endocrinology*.
- Das UB, Mallick M, Debnath JM, Ghosh D (2002) Protective effect of ascorbic acid on cyclophosphamide- induced testicular gametogenic and androgenic disorders in male rats. *Asian J. Androl*, **4**: 201-207.
- Davis BK (1981) Timing of fertilization in mammals: sperm cholesterol/phospholipid ratio as a determinant of the capacitation interval. *Proc. Natl. Acad. Sci. U. S. A.*, **78**: 7560-7564.
- Davis JT, Bridges RB, Coniglio JG (1966) Changes in lipid composition of the maturing rat testis. *Biochem. J.*, **98**: 342-346.
- Davis JT, Coniglio JG (1966) The biosynthesis of docosapentaenoic and other fatty acids by rat testes. *J. Biol. Chem.*, **241**: 610-612.
- Delic JI, Hendry JH, Morris ID, Shalet SM (1986) Dose and time relationships in the endocrine response of the irradiated adult rat testis. *J. Androl*, **7**: 32-41.
- Eckhardt M, Yaghootfam A, Fewou SN, Zoller I, Gieselmann V (2005) A mammalian fatty acid hydroxylase responsible for the formation of alpha-hydroxylated galactosylceramide in myelin. *Biochem. J.*, **388**: 245-254.
- Fan J, Lefebvre J, Manjunath P (2006) Bovine seminal plasma proteins and their relatives: A new expanding superfamily in mammals. *Gene*, **375**: 63-74.
- Flesch FM, Gadella BM (2000) Dynamics of the mammalian sperm plasma membrane in the process of fertilization. *Biochim. Biophys. Acta*, **1469**: 197-235.
- Fofana M, Maboundou JC, Bocquet J, Le GD (1996) Transfer of cholesterol between high density lipoproteins and cultured rat Sertoli cells. *Biochem. Cell Biol.*, **74**: 681-686.
- Fofana M, Travert C, Carreau S, Le GD (2000) Evaluation of cholesteryl ester transfer in the seminiferous tubule cells of immature rats in vivo and in vitro. *J. Reprod. Fertil.*, **118**: 79-83.
- Fraiser LH, Kanekal S, Kehrer JP (1991) Cyclophosphamide toxicity. Characterising and avoiding the problem. *Drugs*, **42**: 781-795.
- Fujimoto H, Tadano-Aritomi K, Tokumasu A, Ito K, Hikita T, Suzuki K, Ishizuka I (2000) Requirement of seminolipid in spermatogenesis revealed by UDP-galactose: Ceramide galactosyltransferase-deficient mice. *J. Biol. Chem.*, **275**: 22623-22626.
- Furland NE (2006) Lípidos con ácidos grasos poliinsaturados de larga y muy larga cadena de las series n-6 y n-3 en el tracto genital de mamíferos. *Tesis Doctoral*
- Furland NE, Maldonado EN, Aresti PA, Avelldano MI (2007a) Changes in lipids containing long- and very long-chain polyunsaturated fatty acids in cryptorchid rat testes. *Biol. Reprod.*, **77**: 181-188.
- Furland NE, Maldonado EN, Avelldano MI (2003) Very long chain PUFA in murine testicular triglycerides and cholesterol esters. *Lipids*, **38**: 73-80.
- Furland NE, Oresti GM, Antollini SS, Venturino A, Maldonado EN, Avelldano MI (2007b) Very long-chain polyunsaturated fatty acids are the major acyl groups of sphingomyelins and ceramides in the head of mammalian spermatozoa. *J. Biol. Chem.*, **282**: 18151-18161.

- Furland NE, Zanetti SR, Oresti GM, Maldonado EN, Avelano MI (2007c) Ceramides and sphingomyelins with high proportions of very long-chain polyunsaturated fatty acids in mammalian germ cells. *J. Biol. Chem.*, **282**: 18141-18150.
- Futerman AH, Riezman H (2005) The ins and outs of sphingolipid synthesis. *Trends Cell Biol.*, **15**: 312-318.
- Futerman AH, Stieger B, Hubbard AL, Pagano RE (1990) Sphingomyelin synthesis in rat liver occurs predominantly at the cis and medial cisternae of the Golgi apparatus. *J. Biol. Chem.*, **265**: 8650-8657.
- Gadella BM (2008) Sperm membrane physiology and relevance for fertilization. *Anim Reprod. Sci.*, **107**: 229-236.
- Gadella BM, Harrison RA (2000) The capacitating agent bicarbonate induces protein kinase A-dependent changes in phospholipid transbilayer behavior in the sperm plasma membrane. *Development*, **127**: 2407-2420.
- Gitlits VM, Toh BH, Loveland KL, Sentry JW (2000) The glycolytic enzyme enolase is present in sperm tail and displays nucleotide-dependent association with microtubules. *Eur. J. Cell Biol.*, **79**: 104-111.
- Gorgas K, Teigler A, Komljenovic D, Just WW (2006) The ether lipid-deficient mouse: tracking down plasmalogen functions. *Biochim. Biophys. Acta*, **1763**: 1511-1526.
- Grassme H, Riethmuller J, Gulbins E (2007) Biological aspects of ceramide-enriched membrane domains. *Prog. Lipid Res.*, **46**: 161-170.
- Grogan WM (1984) Metabolism of arachidonate in rat testis: characterization of 26-30 carbon polyenoic acids. *Lipids*, **19**: 341-346.
- Grogan WM, Huth EG (1983) Biosynthesis of long-chain polyenoic acids from arachidonic acid in cultures of enriched spermatocytes and spermatids from mouse testis. *Lipids*, **18**: 275-284.
- Grogan WM, Lam JW (1982) Fatty acid synthesis in isolated spermatocytes and spermatids of mouse testis. *Lipids*, **17**: 604-611.
- Gunasekar PG, Kumaran B, Govindarajulu P (1991) Role of prolactin on Leydig, Sertoli and germ cellular neutral lipids in bonnet monkeys, *Macaca radiata*. *Endocrinol. Jpn.*, **38**: 1-8.
- Haimovitz-Friedman A, Kolesnick RN, Fuks Z (1997) Ceramide signaling in apoptosis. *Br. Med. Bull.*, **53**: 539-553.
- Hajra AK (1997) Dihydroxyacetone phosphate acyltransferase. *Biochim. Biophys. Acta*, **1348**: 27-34.
- Hajra AK, Das AK (1996) Lipid biosynthesis in peroxisomes. *Ann. N. Y. Acad. Sci.*, **804**: 129-141.
- Hannun YA, Obeid LM (2002) The Ceramide-centric universe of lipid-mediated cell regulation: stress encounters of the lipid kind. *J. Biol. Chem.*, **277**: 25847-25850.
- Hannun YA, Obeid LM (2008) Principles of bioactive lipid signalling: lessons from sphingolipids. *Nat. Rev. Mol. Cell Biol.*, **9**: 139-150.
- Hasegawa M, Wilson G, Russell LD, Meistrich ML (1997) Radiation-induced cell death in the mouse testis: relationship to apoptosis. *Radiat. Res.*, **147**: 457-467.
- Hemminki K, Kallama S (1986) Reactions of nitrogen mustards with DNA. *IARC Sci. Publ.*, 55-70.
- Henkel RR, Schill WB (2003) Sperm preparation for ART. *Reprod. Biol. Endocrinol.*, **1**: 108.
- Henriksen K, Kulmala J, Toppari J, Mehrotra K, Parvinen M (1996) Stage-specific apoptosis in the rat seminiferous epithelium: quantification of irradiation effects. *J. Androl*, **17**: 394-402.
- Hermo L, Chung S, Gregory M, Smith CE, Wang SP, El-Alfy M, Cyr DG, Mitchell GA, Trasler J (2008) Alterations in the testis of hormone sensitive lipase-deficient mice is associated with decreased sperm counts, sperm motility, and fertility. *Mol. Reprod. Dev.*, **75**: 565-577.

- Hinkovska VT, Petkova DH, Koumanov KS (1987) A neutral sphingomyelinase in spermatozoal plasma membranes. *Biochem. Cell Biol.*, **65**: 525-528.
- Hinkovska-Galcheva VT, Petkova DH, Nikolova MN (1989) Sphingomyelin and ceramide: phosphoethanolamine synthesis in ram spermatozoa plasma membrane. *Int. J. Biochem.*, **21**: 1153-1156.
- Holub BJ, Skeaff CM (1987) Nutritional regulation of cellular phosphatidylinositol. *Methods Enzymol.*, **141**: 234-244.
- Hussein MR, bu-Dief EE, bou El-Ghait AT, Adly MA, Abdelraheem MH (2006) Morphological evaluation of the radioprotective effects of melatonin against X-ray-induced early and acute testis damage in Albino rats: an animal model. *Int. J. Exp. Pathol.*, **87**: 237-250.
- Huyghe S, Schmalbruch H, De GK, Verhoeven G, Guillou F, Van Veldhoven PP, Baes M (2006) Peroxisomal multifunctional protein 2 is essential for lipid homeostasis in Sertoli cells and male fertility in mice. *Endocrinology*, **147**: 2228-2236.
- Ishizuka I (1997) Chemistry and functional distribution of sulfoglycolipids. *Prog. Lipid Res.*, **36**: 245-319.
- Ishizuka I, Suzuki M, Yamakawa T (1973) Isolation and characterization of a novel sulfoglycolipid, 'seminolipid,' from boar testis and spermatozoa. *J. Biochem.*, **73**: 77-87.
- Jakobsson A, Westerberg R, Jacobsson A (2006) Fatty acid elongases in mammals: their regulation and roles in metabolism. *Prog. Lipid Res.*, **45**: 237-249.
- Jegou B (1991) Spermatids are regulators of Sertoli cell function. *Ann. N. Y. Acad. Sci.*, **637**: 340-353.
- Kabbaj O, Holm C, Vitale ML, Pelletier RM (2001) Expression, activity, and subcellular localization of testicular hormone-sensitive lipase during postnatal development in the guinea pig. *Biol. Reprod.*, **65**: 601-612.
- Kabbaj O, Yoon SR, Holm C, Rose J, Vitale ML, Pelletier RM (2003) Relationship of the hormone-sensitive lipase-mediated modulation of cholesterol metabolism in individual compartments of the testis to serum pituitary hormone and testosterone concentrations in a seasonal breeder, the mink (*Mustela vison*). *Biol. Reprod.*, **68**: 722-734.
- Kangasniemi M, Huhtaniemi I, Meistrich ML (1996) Failure of spermatogenesis to recover despite the presence of a spermatogonia in the irradiated LBNF1 rat. *Biol. Reprod.*, **54**: 1200-1208.
- Kawasaki Y, Nakagawa A, Nagaosa K, Shiratsuchi A, Nakanishi Y (2002) Phosphatidylserine binding of class B scavenger receptor type I, a phagocytosis receptor of testicular sertoli cells. *J. Biol. Chem.*, **277**: 27559-27566.
- Kerr JB, De Kretser DM (1974) Proceedings: The role of the Sertoli cell in phagocytosis of the residual bodies of spermatids. *J. Reprod. Fertil.*, **36**: 439-440.
- Kerr JB, Mayberry RA, Irby DC (1984) Morphometric studies on lipid inclusions in Sertoli cells during the spermatogenic cycle in the rat. *Cell Tissue Res.*, **236**: 699-709.
- Knobil y Neill's (2006) in: The Physiology of Reproduction, 3rd edition, volume . *Elsevier inc.*
- Kolesnick R, Fuks Z (2003) Radiation and ceramide-induced apoptosis. *Oncogene*, **22**: 5897-5906.
- Kolesnick RN, Goni FM, Alonso A (2000) Compartmentalization of ceramide signaling: physical foundations and biological effects. *J. Cell Physiol*, **184**: 285-300.
- Kolodny EH (2000) Niemann-Pick disease. *Curr. Opin. Hematol.*, **7**: 48-52.
- Kornblatt MJ, Knapp A, Levine M, Schachter H, Murray RK (1974) Studies on the structure and formation during spermatogenesis of the sulfoglycerogalactolipid of rat testis. *Can. J. Biochem.*, **52**: 689-697.
- Larson JL, Miller DJ (1999) Simple histochemical stain for acrosomes on sperm from several species. *Mol. Reprod. Dev.*, **52**: 445-449.

- Leblond E, Desnoyers L, Manjunath P (1993) Phosphorylcholine-binding proteins from the seminal fluids of different species share antigenic determinants with the major proteins of bovine seminal plasma. *Mol. Reprod. Dev.*, **34**: 443-449.
- Lee K, Park JS, Kim YJ, Soo Lee YS, Sook Hwang TS, Kim DJ, Park EM, Park YM (2002) Differential expression of Prx I and II in mouse testis and their up-regulation by radiation. *Biochem. Biophys. Res. Commun.*, **296**: 337-342.
- Lee NP, Mruk DD, Xia W, Cheng CY (2007) Cellular localization of sphingomyelin synthase 2 in the seminiferous epithelium of adult rat testes. *J. Endocrinol.*, **192**: 17-32.
- Lizama C, Alfaro I, Reyes JG, Moreno RD (2007) Up-regulation of CD95 (Apo-1/Fas) is associated with spermatocyte apoptosis during the first round of spermatogenesis in the rat. *Apoptosis.*, **12**: 499-512.
- Luers GH, Thiele S, Schad A, Volkl A, Yokota S, Seitz J (2006) Peroxisomes are present in murine spermatogonia and disappear during the course of spermatogenesis. *Histochem. Cell Biol.*, **125**: 693-703.
- Manda K, Bhatia AL (2003) Pre-administration of beta-carotene protects tissue glutathione and lipid peroxidation status following exposure to gamma radiation. *J. Environ. Biol.*, **24**: 369-372.
- Manda K, Ueno M, Anzai K (2007a) AFMK, a melatonin metabolite, attenuates X-ray-induced oxidative damage to DNA, proteins and lipids in mice. *J. Pineal Res.*, **42**: 386-393.
- Manda K, Ueno M, Moritake T, Anzai K (2007b) alpha-Lipoic acid attenuates x-irradiation-induced oxidative stress in mice. *Cell Biol. Toxicol.*, **23**: 129-137.
- Mandal MN, Ambasadhan R, Wong PW, Gage PJ, Sieving PA, Ayyagari R (2004) Characterization of mouse orthologue of ELOVL4: genomic organization and spatial and temporal expression. *Genomics*, **83**: 626-635.
- Manjunath P, Lefebvre J, Jois PS, Fan J, Wright MW (2008) New Nomenclature for Mammalian BSP Genes. *Biol. Reprod.* .
- Manjunath P, Therien I (2002) Role of seminal plasma phospholipid-binding proteins in sperm membrane lipid modification that occurs during capacitation. *J. Reprod. Immunol.*, **53**: 109-119.
- Martinez P, Morros A (1996) Membrane lipid dynamics during human sperm capacitation. *Front Biosci.*, **1**: d103-d117.
- Marzouki ZM, Coniglio JG (1982) Effect of essential fatty acid deficiency on lipids of rat Sertoli and germinal cells. *Biol. Reprod.*, **27**: 312-315.
- Marzouki ZM, Coniglio JG (1984) The effects of hypophysectomy and testosterone treatment on the composition and metabolism of testicular lipids. *Lipids*, **19**: 609-616.
- Mauduit C, Siah A, Foch M, Chapet O, Clippe S, Gerard JP, Benahmed M (2001) Differential expression of growth factors in irradiated mouse testes. *Int. J. Radiat. Oncol. Biol. Phys.*, **50**: 203-212.
- Megha, Bakht O, London E (2006) Cholesterol precursors stabilize ordinary and ceramide-rich ordered lipid domains (lipid rafts) to different degrees. Implications for the Bloch hypothesis and sterol biosynthesis disorders. *J. Biol. Chem.*, **281**: 21903-21913.
- Megha, London E (2004) Ceramide selectively displaces cholesterol from ordered lipid domains (rafts): implications for lipid raft structure and function. *J. Biol. Chem.*, **279**: 9997-10004.
- Meistrich ML, Hunter NR, Suzuki N, Trostle PK, Withers HR (1978) Gradual regeneration of mouse testicular stem cells after exposure to ionizing radiation. *Radiat. Res.*, **74**: 349-362.
- Meistrich ML, Mohapatra B, Shirley CR, Zhao M (2003) Roles of transition nuclear proteins in spermiogenesis. *Chromosoma*, **111**: 483-488.
- Meistrich ML, Shetty G (2003) Inhibition of spermatogonial differentiation by testosterone. *J. Androl.*, **24**: 135-148.

- Meistrich ML, Wilson G, Shuttlesworth G, Huhtaniemi I, Reissmann T (2001) GnRH agonists and antagonists stimulate recovery of fertility in irradiated LBNF1 rats. *J. Androl*, **22**: 809-817.
- Miller DK (1999) Activation of apoptosis and its inhibition. *Ann. N. Y. Acad. Sci.*, **886**: 132-157.
- Mizutani Y, Kihara A, Chiba H, Tojo H, Igarashi Y (2008) 2-Hydroxy-ceramide synthesis by ceramide synthase family: enzymatic basis for the preference of FA chain length. *J. Lipid Res.*, **49**: 2356-2364.
- Mizutani Y, Kihara A, Igarashi Y (2006) LASS3 (longevity assurance homologue 3) is a mainly testis-specific (dihydro)ceramide synthase with relatively broad substrate specificity. *Biochem. J.*, **398**: 531-538.
- Mohammed BS, Sankarappa S, Geiger M, Sprecher H (1995) Reevaluation of the pathway for the metabolism of 7,10,13, 16-docosatetraenoic acid to 4,7,10,13,16-docosapentaenoic acid in rat liver. *Arch. Biochem. Biophys.*, **317**: 179-184.
- Morris Kates (1972) Technics in lipidology. *Eds. Work TS y Work E. Elsevier pp. 570*
- Murase T, Imaeda N, Kondoh N, Tsubota T (2004) Ceramide enhances acrosomal exocytosis triggered by calcium and the calcium ionophore A23187 in boar spermatozoa. *J. Reprod. Dev.*, **50**: 667-674.
- Nakagawa A, Shiratsuchi A, Tsuda K, Nakanishi Y (2005) In vivo analysis of phagocytosis of apoptotic cells by testicular Sertoli cells. *Mol. Reprod. Dev.*, **71**: 166-177.
- Nakanishi Y, Shiratsuchi A (2004) Phagocytic removal of apoptotic spermatogenic cells by Sertoli cells: mechanisms and consequences. *Biol. Pharm. Bull.*, **27**: 13-16.
- Nenicu A, Luers GH, Kovacs W, David M, Zimmer A, Bergmann M, Baumgart-Vogt E (2007) Peroxisomes in human and mouse testis: differential expression of peroxisomal proteins in germ cells and distinct somatic cell types of the testis. *Biol. Reprod.*, **77**: 1060-1072.
- Nikolopoulou M, Soucek DA, Vary JC (1986) Modulation of the lipid composition of boar sperm plasma membranes during an acrosome reaction in vitro. *Arch. Biochem. Biophys.*, **250**: 30-37.
- Oku H, Shudo J, Mimura K, Haratake A, Nagata J, Chinen I (1995) 1-O-alkyl-2,3-diacylglycerols in the skin surface lipids of the hairless mouse. *Lipids*, **30**: 169-172.
- Osawa Y, Uchinami H, Bielawski J, Schwabe RF, Hannun YA, Brenner DA (2005) Roles for C16-ceramide and sphingosine 1-phosphate in regulating hepatocyte apoptosis in response to tumor necrosis factor- α . *J. Biol. Chem.*, **280**: 27879-27887.
- Osuga J, Ishibashi S, Oka T, Yagyu H, Tozawa R, Fujimoto A, Shionoiri F, Yahagi N, Kraemer FB, Tsutsumi O, Yamada N (2000) Targeted disruption of hormone-sensitive lipase results in male sterility and adipocyte hypertrophy, but not in obesity. *Proc. Natl. Acad. Sci. U. S. A.*, **97**: 787-792.
- Ojala M, Pentikainen MO, Matikainen T, Suomalainen L, Hakala JK, Perez GI, Tenhunen M, Erkkila K, Kovanen P, Parvinen M, Dunkel L (2005) Effects of acid sphingomyelinase deficiency on male germ cell development and programmed cell death. *Biol. Reprod.*, **72**: 86-96.
- Park EM, Park JS, Kim YS, Sung JS, Hwang TS, Kim WC, Han MY, Park YM (2001) Role of oxidative stress in the radiation-induced lung pathogenesis in mice. *J. Biochem. Mol. Biol.* 34-544-550.
- Park HS, Kim S, Lee Y, Choi MS, Choi MU (2006) Alteration of lipid composition of rat thymus during thymic atrophy by whole-body X-irradiation. *Int. J. Radiat. Biol.*, **82**: 129-137.
- Pentikainen V, Dunkel L, Erkkila K (2003) Male germ cell apoptosis. *Endocr. Dev.*, **5**: 56-80.
- Pickering SJ, Fleming TP, Braude PR, Bolton VN, Gresham GA (1989) Are human spermatozoa separated on a Percoll density gradient safe for therapeutic use? *Fertil. Steril.*, **51**: 1024-1029.
- Pineau C, Le MB, Courtens JL, Jegou B (1991) Study in vitro of the phagocytic function of Sertoli cells in the rat. *Cell Tissue Res.*, **264**: 589-598.

- Pinon-Lataillade G, Velez de la Calle JF, Viguier-Martinez MC, Garnier DH, Folliot R, Maas J, Jegou B (1988) Influence of germ cells upon Sertoli cells during continuous low-dose rate gamma-irradiation of adult rats. *Mol. Cell Endocrinol.*, **58**: 51-63.
- Pinon-Lataillade G, Viguier-Martinez MC, Touzalin AM, Maas J, Jegou B (1991) Effect of an acute exposure of rat testes to gamma rays on germ cells and on Sertoli and Leydig cell functions. *Reprod. Nutr. Dev.*, **31**: 617-629.
- Porn MI, Tenhunen J, Slotte JP (1991) Increased steroid hormone secretion in mouse Leydig tumor cells after induction of cholesterol translocation by sphingomyelin degradation. *Biochim. Biophys. Acta*, **1093**: 7-12.
- Porter KL, Shetty G, Meistrich ML (2006) Testicular edema is associated with spermatogonial arrest in irradiated rats. *Endocrinology*, **147**: 1297-1305.
- Poulos A, Johnson DW, Beckman K, White IG, Easton C (1987) Occurrence of unusual molecular species of sphingomyelin containing 28-34-carbon polyenoic fatty acids in ram spermatozoa. *Biochem. J.*, **248**: 961-964.
- Poulos A, Sharp P, Johnson D, White I, Fellenberg A (1986) The occurrence of polyenoic fatty acids with greater than 22 carbon atoms in mammalian spermatozoa. *Biochem. J.*, **240**: 891-895.
- Prabhu SM, Meistrich ML, McLaughlin EA, Roman SD, Warne S, Mendis S, Itman C, Loveland KL (2006) Expression of c-Kit receptor mRNA and protein in the developing, adult and irradiated rodent testis. *Reproduction.*, **131**: 489-499.
- Rabionet M, van der Spoel AC, Chuang CC, von Tumpling-Radosta B, Litjens M, Bouwmeester D, Hellbusch CC, Korner C, Wiegandt H, Gorgas K, Platt FM, Grone HJ, Sandhoff R (2008) Male germ cells require polyenoic sphingolipids with complex glycosylation for completion of meiosis: a link to ceramide synthase-3. *J. Biol. Chem.*, **283**: 13357-13369.
- Ramstedt B, Leppimaki P, Axberg M, Slotte JP (1999) Analysis of natural and synthetic sphingomyelins using high-performance thin-layer chromatography. *Eur. J. Biochem.*, **266**: 997-1002.
- Rana AP, Majumder GC, Misra S, Ghosh A (1992) Occurrence of wax esters and 1-O-alkyl-2,3-diacylglycerols in goat epididymal sperm plasma membrane. *Lipids*, **27**: 75-78.
- Reisse S, Rothardt G, Volkl A, Beier K (2001) Peroxisomes and ether lipid biosynthesis in rat testis and epididymis. *Biol. Reprod.*, **64**: 1689-1694.
- Retterstol K, Tran TN, Haugen TB, Christophersen BO (2001) Metabolism of very long chain polyunsaturated fatty acids in isolated rat germ cells. *Lipids*, **36**: 601-606.
- Reyes JG, Diaz A, Osses N, Opazo C, Benos DJ (1997) On stage single cell identification of rat spermatogenic cells. *Biol. Cell*, **89**: 53-66.
- Reynolds CP, Maurer BJ, Kolesnick RN (2004) Ceramide synthesis and metabolism as a target for cancer therapy. *Cancer Lett.*, **206**: 169-180.
- Robertson KM, Schuster GU, Steffensen KR, Hovatta O, Meaney S, Hultenby K, Johansson LC, Svechnikov K, Soder O, Gustafsson JA (2005) The liver X receptor- β is essential for maintaining cholesterol homeostasis in the testis. *Endocrinology*, **146**: 2519-2530.
- Robinson BS, Johnson DW, Poulos A (1992) Novel molecular species of sphingomyelin containing 2-hydroxylated polyenoic very-long-chain fatty acids in mammalian testes and spermatozoa. *J. Biol. Chem.*, **267**: 1746-1751.
- Rodriguez I, Ody C, Araki K, Garcia I, Vassalli P (1997) An early and massive wave of germinal cell apoptosis is required for the development of functional spermatogenesis. *EMBO J.*, **16**: 2262-2270.
- Roldan ER, Murase T (1994) Polyphosphoinositide-derived diacylglycerol stimulates the hydrolysis of phosphatidylcholine by phospholipase C during exocytosis of the ram sperm acrosome. Effect is not mediated by protein kinase C. *J. Biol. Chem.*, **269**: 23583-23589.

- Roldan ER, Vazquez JM (1996) Bicarbonate/CO₂ induces rapid activation of phospholipase A₂ and renders boar spermatozoa capable of undergoing acrosomal exocytosis in response to progesterone. *FEBS Lett.*, **396**: 227-232.
- Romrell LJ, Bellve AR, Fawcett DW (1976) Separation of mouse spermatogenic cells by sedimentation velocity. A morphological characterization. *Dev. Biol.*, **49**: 119-131.
- Rotstein NP, Pennacchiotti GL, Sprecher H, Aveldano MI (1996) Active synthesis of C₂₄:5, n-3 fatty acid in retina. *Biochem. J.*, **316 (Pt 3)**: 859-864.
- Rouser G, Fkeischer S, Yamamoto A (1970) Two dimensional thin layer chromatographic separation of polar lipids and determination of phospholipids by phosphorus analysis of spots. *Lipids*, **5**: 494-496.
- Russell LD, de Franca LR (1995) Building a testis. *Tissue Cell*, **27**: 129-147.
- Russell LD, Ettlin RA, Hikim AP, Clegg ED (1990) in: Histological and Histopathological Evaluation of the Testis. *Cache River, Clearwater*
- Saraste A, Pulkki K (2000) Morphologic and biochemical hallmarks of apoptosis. *Cardiovasc. Res.*, **45**: 528-537.
- Selva DM, Hirsch-Reinshagen V, Burgess B, Zhou S, Chan J, McIsaac S, Hayden MR, Hammond GL, Vogl AW, Wellington CL (2004) The ATP-binding cassette transporter 1 mediates lipid efflux from Sertoli cells and influences male fertility. *J. Lipid Res.*, **45**: 1040-1050.
- Selvakumar E, Prahalathan C, Sudharsan PT, Varalakshmi P (2006) Protective effect of lipoic acid on cyclophosphamide-induced testicular toxicity. *Clin. Chim. Acta*, **367**: 114-119.
- Setchell BP, Main SJ (1975) The blood - testis barrier and steroids. *Curr. Top. Mol. Endocrinol.*, **2**: 223-233.
- Shetty G, Meistrich ML (2005) Hormonal approaches to preservation and restoration of male fertility after cancer treatment. *J. Natl. Cancer Inst. Monogr*, 36-39.
- Shetty G, Weng CC, Bolden-Tiller OU, Huhtaniemi I, Handelsman DJ, Meistrich ML (2004) Effects of medroxyprogesterone and estradiol on the recovery of spermatogenesis in irradiated rats. *Endocrinology*, **145**: 4461-4469.
- Shiratsuchi A, Kawasaki Y, Ikemoto M, Arai H, Nakanishi Y (1999) Role of class B scavenger receptor type I in phagocytosis of apoptotic rat spermatogenic cells by Sertoli cells. *J. Biol. Chem.*, **274**: 5901-5908.
- Sikka SC (2001) Relative impact of oxidative stress on male reproductive function. *Curr. Med. Chem.*, **8**: 851-862.
- Sinha Hikim AP, Lue Y, az-Romero M, Yen PH, Wang C, Swerdloff RS (2003) Deciphering the pathways of germ cell apoptosis in the testis. *J. Steroid Biochem. Mol. Biol.*, **85**: 175-182.
- Sleight SB, Miranda PV, Plaskett NW, Maier B, Lysiak J, Scrable H, Herr JC, Visconti PE (2005) Isolation and proteomic analysis of mouse sperm detergent-resistant membrane fractions: evidence for dissociation of lipid rafts during capacitation. *Biol. Reprod.*, **73**: 721-729.
- Smith RE, Lespi P, Di LM, Bustos C, Marra FA, de Alaniz MJ, Marra CA (2008) A reliable biomarker derived from plasmalogens to evaluate malignancy and metastatic capacity of human cancers. *Lipids*, **43**: 79-89.
- Snyder F (1999) The ether lipid trail: a historical perspective. *Biochim. Biophys. Acta*, **1436**: 265-278.
- Songthaveesin C, Saikhun J, Kitiyanant Y, Pavasuthipaisit K (2004) Radio-protective effect of vitamin E on spermatogenesis in mice exposed to gamma-irradiation: a flow cytometric study. *Asian J. Androl*, **6**: 331-336.
- Spence MW, Burgess JK, Sperker ER (1979) Neutral and acid sphingomyelinases: somatotopographical distribution in human brain and distribution in rat organs. A possible relationship with the dopamine system. *Brain Res.*, **168**: 543-551.

- Sulkowska M, Sulkowski S, Skrzydlewska E, Farbiszewski R (1998) Cyclophosphamide-induced generation of reactive oxygen species. Comparison with morphological changes in type II alveolar epithelial cells and lung capillaries. *Exp. Toxicol. Pathol.*, **50**: 209-220.
- Suomalainen L, Hakala JK, Pentikainen V, Ojala M, Erkkila K, Pentikainen MO, Dunkel L (2003) Sphingosine-1-phosphate in inhibition of male germ cell apoptosis in the human testis. *J. Clin. Endocrinol. Metab.*, **88**: 5572-5579.
- Tadano-Aritomi K, Matsuda J, Fujimoto H, Suzuki K, Ishizuka I (2003) Seminolipid and its precursor/degradative product, galactosylalkylacylglycerol, in the testis of saposin A- and prosaposin-deficient mice. *J. Lipid Res.*, **44**: 1737-1743.
- Tafesse FG, Huitema K, Hermansson M, van der PS, van den DJ, Uphoff A, Somerharju P, Holthuis JC (2007b) Both sphingomyelin synthases SMS1 and SMS2 are required for sphingomyelin homeostasis and growth in human HeLa cells. *J. Biol. Chem.*, **282**: 17537-17547.
- Tafesse FG, Ternes P, Holthuis JC (2006a) The multigenic sphingomyelin synthase family. *J. Biol. Chem.*, **281**: 29421-29425.
- Tanphaichitr N, Agulnick A, Seibel M, Taymor M (1988) Comparison of the in vitro fertilization rate by human sperm capacitated by multiple-tube swim-up and Percoll gradient centrifugation. *J. In Vitro Fert. Embryo. Transf.*, **5**: 119-122.
- Therien I, Bousquet D, Manjunath P (2001) Effect of seminal phospholipid-binding proteins and follicular fluid on bovine sperm capacitation. *Biol. Reprod.*, **65**: 41-51.
- Tripathi DN, Jena GB (2008) Astaxanthin inhibits cytotoxic and genotoxic effects of cyclophosphamide in mice germ cells. *Toxicology*, **248**: 96-103.
- Tvrzicka E, Rezanka T, Krijt J, Janousek V (1988) Identification of very-long-chain fatty acids in rat and mouse harderian gland lipids by capillary gas chromatography-mass spectrometry. *J. Chromatogr.*, **431**: 231-238.
- Uchida Y, Hama H, Alderson NL, Douangpanya S, Wang Y, Crumrine DA, Elias PM, Holleran WM (2007) Fatty acid 2-hydroxylase, encoded by FA2H, accounts for differentiation-associated increase in 2-OH ceramides during keratinocyte differentiation. *J. Biol. Chem.*, **282**: 13211-13219.
- Van Gestel RA, Brewis IA, Ashton PR, Helms JB, Brouwers JF, Gadella BM (2005) Capacitation-dependent concentration of lipid rafts in the apical ridge head area of porcine sperm cells. *Mol. Hum. Reprod.*, **11**: 583-590.
- Velamakanni S, Wei SL, Janvilisri T, van Veen HW (2007) ABCG transporters: structure, substrate specificities and physiological roles : a brief overview. *J. Bioenerg. Biomembr.*, **39**: 465-471.
- Vergouwen RP, Huiskamp R, Bas RJ, Roepers-Gajadien HL, Davids JA, de Rooij DG (1995) Radiosensitivity of testicular cells in the fetal mouse. *Radiat. Res.*, **141**: 66-73.
- Volle DH, Mouzat K, Duggavathi R, Siddeek B, Dechelotte P, Sion B, Veyssiere G, Benahmed M, Lobaccaro JM (2007) Multiple roles of the nuclear receptors for oxysterols liver X receptor to maintain male fertility. *Mol. Endocrinol.*, **21**: 1014-1027.
- Voss A, Reinhart M, Sankarappa S, Sprecher H (1991) The metabolism of 7,10,13,16,19-docosapentaenoic acid to 4,7,10,13,16,19-docosahexaenoic acid in rat liver is independent of a 4-desaturase. *J. Biol. Chem.*, **266**: 19995-20000.
- Walburg HE (1975) Radiation-induced life-shortening and premature aging, in: *J.T. Lett, D.H. Adler (Eds.), Advances in Radiation Biology, vol. 5, Academic Press, New York*, pp. 145-179.
- Wang N, Tall AR (2003) Regulation and mechanisms of ATP-binding cassette transporter A1-mediated cellular cholesterol efflux. *Arterioscler. Thromb. Vasc. Biol.*, **23**: 1178-1184.
- Xiong W, Chen Y, Wang H, Wang H, Wu H, Lu Q, Han D (2008) Gas6 and the Tyro 3 receptor tyrosine kinase subfamily regulate the phagocytic function of Sertoli cells. *Reproduction.*, **135**: 77-87.

- Yefimova MG, Sow A, Fontaine I, Guilleminot V, Martinat N, Crepieux P, Canepa S, Maurel MC, Fouhecourt S, Reiter E, Benzakour O, Guillou F (2008) Dimeric transferrin inhibits phagocytosis of residual bodies by testicular rat Sertoli cells. *Biol. Reprod.*, **78**: 697-704.
- Yu C, Alterman M, Dobrowsky RT (2005) Ceramide displaces cholesterol from lipid rafts and decreases the association of the cholesterol binding protein caveolin-1. *J. Lipid Res.*, **46**: 1678-1691.
- Zanetti SR (2008) Lípidos con ácidos grasos poliinsaturados de muy larga cadena normales e hidroxilados en células germinales y espermatozoides. *Tesis Doctoral*
- Zanetti SR, Maldonado EN, Avelano MI (2007) Doxorubicin affects testicular lipids with long-chain (C18-C22) and very long-chain (C24-C32) polyunsaturated fatty acids. *Cancer Res.*, **67**: 6973-6980.
- Zhang Y, Hayashi Y, Cheng X, Watanabe T, Wang X, Taniguchi N, Honke K (2005) Testis-specific sulfoglycolipid, seminolipid, is essential for germ cell function in spermatogenesis. *Glycobiology*, **15**: 649-654.
- Zhang Z, Shao S, Meistrich ML (2007) The radiation-induced block in spermatogonial differentiation is due to damage to the somatic environment, not the germ cells. *J. Cell Physiol*, **211**: 149-158.