

Abbracchio MP, Burnstock G. Purinoceptors: are there families of P2X and P2Y purinoceptors? *Pharmacol Ther.* 64: 445-475, 1994.

Abbracchio MP, Burnstock G. Purinergic signaling: pathophysiological roles. *Jpn J Pharmacol.* 78: 113-145, 1998.

Abbracchio MP, Burnstock G, Boeynaems J-M, Barnard EA, Boyer JL, Kennedy C, Knight GE, Fumagalli M, Gachet C, Jacobson KA, Weisman GA. International Union of Pharmacology LVIII: Update on the P2Y G protein-coupled nucleotide receptors: from molecular mechanisms and pathophysiology to therapy. *Pharmacol Rev.* 58: 281-341, 2006.

Abbracchio MP, Ceruti S. Roles of P2 receptors in glial cells: focus on astrocytes. *Purinergic Signal.* 2: 595-604, 2006.

Amaravadi R, Thompson CB. The survival kinases Akt and Pim as potential pharmacological targets. *J Clin Invest.* 115: 2618-2624, 2005.

Baldi C, Vazquez G, Boland R. Characterization of a 1,25(OH)₂-vitamin D₃-responsive capacitance Ca²⁺ entry pathway in rat osteoblast-like cells. *J Cell Biochem.* 86: 678-87, 2002.

Balendran A, Casamayor A, Deak M, Paterson A, Gaffney P, Currie R, Downes CP, Alessi DR. PDK1 acquires PDK2 activity in the presence of a synthetic peptide derived from the carboxyl terminus of PRK2. *Curr Biol.* 9: 393-404, 1999.

Ballermann BJ, Dardik A, Eng E, Liu A. Shear stress and the endothelium. *Kidney Int Suppl.* 67: S100-S108, 1998.

Bard F, Mazelin L, Pechoux-Longin C, Malhotra V, Jurdic P. Src regulates Golgi structure and KDEL receptor-dependent retrograde transport to the endoplasmic reticulum. *J Biol Chem.* 278: 46601-46606, 2003.

Benjamin M, Hiller B. Mechanical influences on cells, tissues and organs - 'Mechanical Morphogenesis'. *Eur J Morphol.* 41: 3-7, 2003.

Bennett DW, Drury AN. Further observations related to the physiological activity of adenine compounds. *J Physiol.* 72: 288-320, 1931.

Biscardi JS, Ishizawa RC, Silva CM, Parsons SJ. Tyrosine kinase signaling in breast cancer: epidermal growth factor receptor and c-Src interactions in breast cancer. *Breast Cancer Res.* 2: 203-210, 2000.

Blanco-Aparicio C, Renner O, Leal JF, Carnero A. PTEN, more than the AKT pathway. *Carcinogenesis*. 28: 1379-1386, 2007.

Bleasdale JE, Thaker R, Gremban RS, Bundy GL, Fitzpatrick FA, Smith RJ. Selective inhibition of receptor-coupled phospholipase C-dependent processes in human platelets and polymorphonuclear neutrophils. *J Pharmacol Exp Ther*. 255, 756-768, 1990.

Bodin P, Burnstock G. Purinergic signaling: ATP release. *Neurochem Res*. 26: 959-969, 2001.

Bowler WB, Dixon CJ, Halleux C, Maier R, Bilbe G, Fraser W, Gallagher J, Hipskind R. Signaling in human osteoblasts by extracellular nucleotides. Their weak induction of the c-fos proto-oncogene via Ca^{2+} mobilization is strongly potentiated by a parathyroid hormone/cAMP-dependent protein kinase pathway independently of mitogen-activated protein kinase. *J Biol Chem*. 274: 14315-14324, 1999.

Bradford M. A rapid and sensitive method for quantification of microgram quantities of proteins utilizing the principle of protein binding. *Anal Biochem*. 72: 248-254, 1976.

Brandt DT, Goerke A, Heuer M, Gimona M, Leitgs M, Kremmer E, Lammers R, Haller H, Mischak H. Protein kinase C delta induces Src kinase activity via activation of the Protein Tyrosine Phosphatase PTP α . *J Biol Chem*. 278: 34073-34078, 2003.

Brautigam V, Frasier C, Nikodemova M, Watters JJ. Purinergic receptor modulation of BV-2 microglial cell activity: potential involvement of p38 MAP kinase and CREB. *J Neuroimmunol*. 166: 113-125, 2005.

Brazil DP, Hemmings BA. Ten years of protein kinase B signaling: a hard Akt to follow. *Trends Biochem Sci*. 26: 657-664, 2001.

Briata P, Forcales SV, Ponassi M, Corte G, Chen CY, Karin M, Puri PL, Gherzi R. p38-dependent phosphorylation of the mRNA decay-promoting factor KSRP controls the stability of select myogenic transcripts. *Mol Cell*. 20: 891-903, 2005.

Burnstock G. Purinergic nerves. *Pharmacol Rev*. 24: 509-581, 1972.

Burnstock G, Kennedy C. Is there a basis for distinguishing two types of P2-purinoceptor? *Gen Pharmacol*. 16: 433-440, 1985.

- Burnstock G. Purinergic signaling. *Br J Pharmacol.* 147: S172-S181, 2006.
- Burnstock G. Purine and pyrimidine receptors. *Cell Mol Life Sci.* 64: 1471-1483, 2007.
- Chaban VV, McRoberts JA, Ennes HS, Mayer EA. Nitric oxide synthase inhibitors enhance mechanosensitive Ca(2+) influx in cultured dorsal root ganglion neurons. *Brain Res.* 903: 74-85, 2001.
- Chang SJ, Tzeng CR, Lee YH, Tai CJ. Extracellular ATP activates the PLC/PKC/ERK signaling pathway through the P2Y2 purinergic receptor leading to the induction of early growth response 1 expression and the inhibition of viability in human endometrial stromal cells. *Cell Signal.* 20: 1248-1255, 2008.
- Chen Y, Sun A. Activation of transcription factor AP-1 by extracellular ATP in PC12 cells. *Neurochem Res.* 23: 543-550, 1998.
- Choudhury GG, Mahimainathan L, Das F, Venkatesan B, Ghosh-Choudhury. *Cell Signal.* 18: 1854-1864, 2006.
- Chung H, Seo S, Moon M, Park S. Phosphatidylinositol-3-kinase/Akt/glycogen synthase kinase-3{beta} and ERK1/2 pathways mediate protective effects of acylated and unacylated ghrelin against oxygen-glucose deprivation-induced apoptosis in primary rat cortical neuronal cells. *J Endocrinol.* Trabajo en prensa, 2008.
- Ciccarelli M, Cipolletta E, Santulli G, Campanile A, Pumiglia K, Cervero P, Pastore L, Astone D, Trimarco B, Iaccarino G. Endothelial beta(2) adrenergic signaling to Akt: Role of G(i) and Src. *Cell Signal.* 19: 1949-1955, 2007.
- Coutinho-Silva R, Stahl L, Cheung K-K, Enes de Campos N, Oliveira Souza C, Ojcius DM, Burnstock G. P2X and P2Y purinergic receptors on human intestinal epithelial cells: effects of extracellular nucleotides on apoptosis and cell proliferation. *Am J Physiol Gastrointest Liver Physiol.* 288: G1024-G1035, 2005.
- Cuenda A, Rousseau S. p38 MAP-kinases pathway regulation, function and role in human diseases. *Bioch Bioph Acta.* 1773: 1358-1375, 2007.
- Cuevas BD, Lu Y, Mao M, Zhang J, LaPushin R, Siminovitch K, Mills G B. Tyrosine Phosphorylation of p85 Relieves Its Inhibitory Activity on Phosphatidylinositol 3-Kinase. *J Biol Chem.* 276: 27455-27461, 2001.

- David-Pfeuty T, Bagrodia S, Shalloway D. Differential localization patterns of myristoylated and nonmyristoylated c-Src proteins in interphase and mitotic c-Src overexpresser cells. *J Cell Sci.* 105, 613-628, 1993.
- Dawes NJ, Cox VM, Park KS, Nga H, Goldspink DF. The induction of c-fos and c-jun in stretched latissimus dorsi muscle of the rabbit: responses to duration, degree and re-application of the stretch stimulus. *Exp Physiol.* 81: 329-339, 1996.
- De Cesare D, Fimia GM, Sassone-Corsi P. Signaling routes to CREM and CREB: plasticity in transcriptional activation. *TIBS.* 24: 281-285, 1999.
- Della Fazia MA, Servillo G, Sassone-Corsi P. Cyclic AMP signaling and cellular proliferation: regulation of CREB and CREM. *FEBS Lett.* 410: 22-24, 1997.
- Di Virgilio F, Chiozzi P, Ferrari D, Falzoni S, Sanz JM, Morelli A, Torboli M, Bolognesi G, Baricordi OR. Nucleotide receptors: an emerging family of regulatory molecules in blood cells. *Blood.* 97: 587-600, 2001.
- Dixon CJ, Bowler WB, Fleetwood P, Ginty AF, Gallagher JA, Carron JA. Extracellular nucleotides stimulate proliferation in MCF-7 breast cancer cells via P2-purinoceptors. *Br J Cancer.* 75: 34-39, 1997.
- Dreskin SC, Thomas GW, Dale SN, Heasley LE. Isoforms of Jun Kinase are differentially expressed and activated in human monocyte/macrophage (THP-1) cells. *J Immunol.* 166: 5646-5653, 2001.
- Drury AN, Szent-Györgyi AY. The physiological activity of adenine compounds with special reference to their action upon the mammalian heart. *J Physiol.* 68: 213-237, 1929.
- Elia MG, Muscella A, Romano S, Greco S, Di Jeso B, Verri T, Storelli C, Marsigliante S. Effects of extracellular nucleotides in the thyroid: P2Y2 receptor-mediated ERK1/2 activation and c-Fos induction in PC Cl3 cells. *Cell Signal.* 17: 739-749, 2005.
- Erb L, Liao Z, Seye CI, Weisman GA. P2 receptors: intracellular signalling. *Eur. J. Physiol.* 452: 552-562, 2006.
- Farooq A, Zhou M-M. Structure and regulation of MAPK phosphatases. *Cell Signal.* 16: 769-779, 2004.

Fries JE, Goczalik IM, Wheeler-Schilling TH, Kohler K, Guenther E, Wolf S, Wiedemann PI, Bringmann A, Reichenbach A, Francke M, Pannicke T. Identification of P2Y Receptor Subtypes in Human Müller Glial Cells by Physiology, Single Cell RT-PCR and Immunohistochemistry. *IOVS*. 46: 3000-3007, 2005.

Furuya K, Akita K, Sokabe M. Extracellular ATP mediated mechano-signaling in mammary glands. *Nippon Yakurigaku Zasshi* 123: 397-402, 2004.

Gerasimovskaya EV, Tucker DA, Weiser-Evans M, Wenzlau JM, Klemm DJ, Banks M, Stenmark KR. Extracellular ATP-induced proliferation of adventitial fibroblasts requires phosphoinositide 3-kinase, Akt, mammalian target of rapamycin and p70 S6 kinase signaling pathways. *J Biol Chem*. 280: 1838-1848, 2005.

Gerasimovskaya EV, Woodward HN, Tucker DA, Stenmark KR. Extracellular ATP is a pro-angiogenic factor for pulmonary artery vasa vasorum endothelial cells. *Angiogenesis* 11: 169-182, 2008.

Grynkiewicz G, Poenie M, Tsien RY. A new generation of Ca^{2+} indicators with greatly improved fluorescence properties. *J Biol Chem*. 260: 3440-3450, 1985.

Go YM, Park H, Maland MC, Darley-Usmar VM, Stoyanov B, Wetzker R, Jo H. Phosphatidylinositol 3-kinase mediates shear stress-dependent activation of JNK in endothelial cells. *Am J Physiol Heart Circ Physiol*. 275: 1898-1904, 1998.

Hanada M, Feng J, Hemmings BA. Structure, regulation and function of PKB/Akt- a major therapeutic target. *Bioch Biophys Acta*. 1697: 3-16, 2004.

Hart S, Fischer OM, Prenzel N, Zwick-Wallasch E, Schneider M, Hennighausen L, Ullrich A. GPCR-induced migration of breast carcinoma cells depends on both EGFR signal transactivation and EGFR-independent pathways. *Biol Chem*. 386: 845-55, 2005.

Haynes MP, Li L, Sinha D, Russell KS, Hisamoto K, Baron R, Collinge M, Sessa WC, Bender JR. Src Kinase Mediates Phosphatidylinositol 3-Kinase/Akt-dependent Rapid Endothelial Nitric-oxide Synthase Activation by Estrogen. *J Biol Chem*. 278: 2118-2123, 2003.

Heo JS, Han HJ. ATP stimulates mouse embryonic stem cell proliferation via PKC, PI3K/Akt and MAPKs signaling pathways. *Stem cells*. 24: 2637-2648, 2006.

Hildebrandt JP, Plant TD, Meves H. The effects of bradykinin on K⁺ currents in NG108-15 cells treated with U73122, a phospholipase C inhibitor, or neomycin. *Br J Pharmacol*. 120: 841-850, 1997.

Hirose Y, Katayama M, Stokoe D, Haas-Kogan DA, Berger MS, Pieper RO. The p38 mitogen-activated protein kinase pathway links the DNA mismatch repair system to the G2 checkpoint and to resistance to chemotherapeutic DNA-methylating agents. *Mol Cell Biol*. 23: 8306-8315, 2003.

Hook SS, Means AR. Ca²⁺/CaM-dependent kinases: from activation to function. *Annu. Rev. Pharmacol Toxicol*. 41: 471-505, 2001.

Huwiler A, RoÈ Iz W, Dorsch S, Ren S, PfeilschifterHuwiler J. Extracellular ATP and UTP activate the protein kinase B/Akt cascade via the P2Y2 purinoceptor in renal mesangial cells. *Br J Pharmacol*. 136: 520-529, 2002.

Ito K, Caramori G, Adcock IM. Therapeutic potential of phosphatidylinositol 3-kinase inhibitors in inflammatory respiratory disease. *J Pharmacol Exp Ther*. 321: 1-8, 2007.

Jacques-Silva MC, Bernardi A, Rodnight R, Lenz G. ERK, PKC and PI3K/Akt pathways mediate extracellular ATP and adenosine-induced proliferation of U138-MG human glioma cell line. *Oncology*. 67: 450-459, 2004a.

Jacques-Silva MC, Rodnight R, Lenz G, Liao Z, Kong Q, Tran M, Kang Y, Gonzalez FA, Weisman GA, Neary JT. P2X7 receptors stimulate Akt phosphorylation in astrocytes. *Br J Pharmacol*. 141: 1106-1117, 2004b.

Jin W, Yun C, Hobbie A, Martin MJ, Sorensen PH, Kim SJ. Cellular transformation and activation of the phosphoinositide-3-kinase-Akt cascade by the ETV6-NTRK3 chimeric tyrosine kinase requires c-Src. *Cancer Res*. 67: 3192-3200, 2007.

Johnson FM, Gallick GE. Src family nonreceptor tyrosine kinases as molecular targets for cancer therapy. *Anticancer Agents Med. Chem*. 7: 651-659, 2007.

Karin M. The regulation of AP-1 activity by mitogen-activated protein kinases. *J Biol Chem*. 270: 16483-16486, 1995.

Katz M, Amit I, Yarden Y. Regulation of MAPKs by growth factors and receptor tyrosine kinases. *Bioch Bioph Acta*. 1773: 1161-1176, 2007.

Katz S, Boland R, Santillán G. Modulation of ERK1/2 and p38 MAPK signaling pathways by ATP in osteoblasts: Involvement of mechanical stress-activated calcium influx, PKC and Src activation. *Int. J. Biochem. Cell Biol.* 38: 2082-2091, 2006.

Katz S, Boland R, Santillán G. Purinergic (ATP) signaling stimulates JNK1 but not JNK2 MAPK in osteoblast-like cells: Contribution of intracellular Ca(2+) release, stress activated and L-voltage-dependent calcium influx, PKC and Src kinases. *Arch Biochem Biophys.* Trabajo en prensa, 2008.

Kawata A, Mikuni-Takagaki Y. Mechanotransduction in stretched osteocytes-temporal expression of immediate early and other genes. *Biochem Biophys Res Commun.* 246: 404-408, 1998.

Kheifets V, Mochly-Rosen D. Insight into intra- and inter-molecular interactions of PKC: Design of specific modulators of kinase function. *Pharmacol Res.* 55: 467-476, 2007.

Kidd LB, Schabbauer GA, Luyendyk JP, Holscher TD, Tilley RE, Tencati M, Mackman N. Insulin activation of the phosphatidylinositol 3-kinase/protein kinase B (Akt) pathway reduces lipopolysaccharide-induced inflammation in mice. *J Pharmacol Exp Ther.* 326: 348-353, 2008.

Kim GY, Mercer SE, Ewton DZ, Yan Z, Jin K, Friedman E. The stress-activated protein kinases p38 alpha and JNK1 stabilize p21(Cip1) by phosphorylation. *J Biol Chem.* 277: 29792-29802, 2002.

Klein-Nulend J, van der Plas A, Semeins CM, Ajubi NE, Frangos JA, Nijweide PJ, Burger EH. Sensitivity of osteocytes to biomechanical stress in vitro. *FASEB J.* 9: 441-445, 1995.

Krasilnikov MA. Phosphatidylinositol-3 kinase dependent pathways: the role in control of cell growth, survival, and malignant transformation. *Biochemistry (Mosc)* 65: 59-67, 2000.

Kumagai H, Sakamoto H, Guggino S, Filburn CR, Sacktor B. Neurotransmitter regulation of cytosolic calcium in osteoblast-like bone cells. *Calcif Tissue Int.* 45: 251-254, 1989.

Kumahashi N, Ochi M, Kataoka H, Uchio Y, Kakimaru H, Sugawara K, Enomoto K. Involvement of ATP, increase of intracellular calcium and the early expression of c-fos in the repair of rat fetal articular cartilage. *Cell Tissue Res.* 317: 117-128, 2004.

Kumar A, Carrera AC. New functions for PI3K in the control of cell division. *Cell Cycle.* 6: 1696-1698, 2007.

- Kushida N, Kabuyama Y, Yamaguchi O, Homma Y. Essential role for extracellular Ca(2+) in JNK activation by mechanical stretch in bladder smooth muscle cells. *Am J Physiol Cell Physiol*. 281: C1165-C1172, 2001.
- Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature*. 227: 680-685, 1970.
- Lahne M, Gale JE. Damage-induced activation of ERK1/2 in cochlear supporting cells is a hair cell death-promoting signal that depends on extracellular ATP and calcium. *J Neurosci*. 28: 4918-4928, 2008.
- Landschulz WH, Jonson PF, McKnight SL. The leucine zipper: a hypothetical structure common to a new class of DNA binding proteins. *Science*. 240: 1759-1764, 1988.
- Lawrence M, Jivan A, Shao C, Duan L, Goad D, Zaganjor E, Osborne J, McGlynn K, Stippec S, Earnest S, Chen W, Cobb M. The roles of MAPKs in disease. *Cell Research*. 18: 436-442, 2008.
- Lehninger, A.L.; Nelson, D.L.; Cox, M.M. Principles of Biochemistry (4th Ed.) W.H. Freeman and Company, New York. 2005.
- Lehoux S, Castier Y, Tedgui A. Molecular mechanisms of the vascular responses to haemodynamic forces. *J Int Med*. 259: 381-392, 2006.
- Li YS, Haga JH, Chien S. Molecular basis of the effects of shear stress on vascular endothelial cells. *J Biomech*. 38: 1949-1971, 2005.
- Liang K, Jin W, Knuefermann C, Schmidt M, Mills GB, Ang KK, Milas L, Fan Z. Targeting the phosphatidylinositol 3-kinase/Akt pathway for enhancing breast cancer cells to radiotherapy. *Mol Cancer Ther*. 2: 353-360, 2003.
- Liang K, Qin T, Yang Z. Liang K, Qin T, Yang Z. Recent researches on the responses of fibroblasts exposed to cyclic mechanical stretching in vitro. *Sheng Wu Yi Xue Gong Cheng Xue Za Zhi*. 21: 1047-1050, 2004.
- Liu H, Adler AS, Segal E, Chang HY. Transcriptional program mediating entry into cellular quiescence. *PLoS Genetics*, 3: 1-13, 2007.
- Liu J, Liao Z, Camden J, Griffin KD, Garrad RC, Santiago-Pérez LI, González FA, Seye CI, Weisman GA, Erb L. Src homology 3 binding sites in the P2Y2 nucleotide receptor interact with Src and regulate

activities of Src, proline-rich tyrosine kinase 2, and growth factor receptors. *J Biol Chem.* 279: 8212-8218, 2004.

Liu Y, Ludes-Meyers J, Zhang Y, Munoz-Medellin D, Kim HT, Lu C, Ge G, Schiff R, Hilsenbeck S, Osborne CK, Brown PH. Inhibition of AP-1 transcription factor causes blockade of multiple signal transduction pathways and inhibits breast cancer growth. *Oncogene* 21: 7680-7689, 2002.

Lu C, Shen Q, DuPre E, Kim H, Hilsenbeck S, Brown PH. c-Fos is critical for MCF-7 breast cancer cell growth. *Oncogene* 24:6516-6524, 2005.

Mackay HJ, Twelves CJ. Protein kinase C: a target for anticancer drugs? *Endocr Relat Canc.* 10: 389-396, 2003.

Marqués M, Kumar A, Cortés I, Gonzalez-García A, Hernández C, Moreno-Ortiz MC, Carrera AC. Phosphoinositide 3-kinases p110alpha and p110beta regulate cell cycle entry, exhibiting distinct activation kinetics in G1 phase. *Mol Cell Biol.* 28: 2803-2814, 2008.

Maruyama T, Kanaji T, Nakade S, Kanno T, Mikoshiba K. 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P₃-induced Ca²⁺ release. *J Biochem (Tokyo).* 122: 498-505, 1997.

Migita K, Lu L, Zhao Y, Honda K, Iwamoto T, Kita S, Katsuragi T. Adenosine induces ATP release via an inositol 1,4,5-trisphosphate signaling pathway in MDCK cells. *Biochem Biophys Res Commun.* 328: 1211-1215, 2005.

Monje P, Marinissen MJ, Gutkind JS. Phosphorylation of the carboxyl-terminal transactivation domain of c-Fos by extracellular signal-regulated kinase mediates the transcriptional activation of AP-1 and cellular transformation induced by platelet-derived growth factor. *Mol Cell Biol.* 23: 7030-7043, 2003.

Montiel M, Pérez de la Blanca E, Jiménez E. P2Y Receptors Activate MAPK/ERK through a Pathway Involving PI3K/PDK1/PKC- ζ in Human Vein Endothelial Cells. *Cell Physiol Biochem.* 18: 123-134, 2006.

Muscella A, Elia MG, Greco S, Storelli C, Marsigliante S. Activation of P2Y₂ receptor induces c-FOS protein through a pathway involving mitogen - activated protein kinases and phosphoinositide 3-kinases in HeLa cells. *J Cell Physiol.* 195: 234-40, 2003.

Naemsch LN, Dixon SJ, Sims SM. Activity-dependent development of P2X7 current and Ca²⁺ entry in rabbit osteoclasts. *J Biol Chem.* 276: 39107-39114, 2001.

Nandigama R, Padmasekar M, Wartenberg M, Saber H. Feed forward cycle of hypotonic stress-induced ATP release, purinergic receptor activation, and growth stimulation of prostate cancer cells. *J Biol Chem.* 281: 5686-5693, 2006.

Neary JT. MAPK cascades in cell growth and death. *News Physiol Sci.* 12: 286-293, 1997.

Neary JT, Kang Y, Bu Y, Yu E, Akong K, Peters CM. Mitogenic signaling by ATP/P2Y purinergic receptors in astrocytes: involvement of a calcium-independent protein kinase C, extracellular signal-regulated protein kinase pathway distinct from the phosphatidylinositol-specific phospholipase C/calcium pathway. *J Neurosci.* 19: 4211-4220, 1999.

Neary JT, Whitemore SR, Bu Y, Mehta H, Shi YF. Biochemical mechanisms of action of Hypericum LI 160 in glial and neuronal cells: inhibition of neurotransmitter uptake and stimulation of extracellular signal regulated protein kinase. *Pharmacopsychiatry.* 34: S103-S107, 2001.

Neary JT, Kang Y, Willoughby KA, Ellis EF. Activation of extracellular signal-regulated kinase by stretch-induced injury in astrocytes involves extracellular ATP and P2 purinergic receptors. *J Neurosci.* 23: 2348-2356, 2003.

Neary JT, Kang Y, Shi YF. Signaling from nucleotide receptors to protein kinase cascades in astrocytes. *Neurochem Res.* 29: 2037-2042, 2004.

Neary JT, Kang Y, Tran M, Feld J. Traumatic injury activates protein kinase B/Akt in cultured astrocytes: role of extracellular ATP and P2 purinergic receptors. *J Neurotrauma.* 22: 491-500, 2005.

Numata T, Yoshino M. Characterization of stretch-activated calcium permeable cation channels in freshly isolated myocytes of the cricket (*Gryllus bimaculatus*) lateral oviduct. *J Insect Physiol.* 51:481-8, 2005.

Ohsawa K, Irino Y, Nakamura Y, Akazawa C, Inoue K, Kohsaka S. Involvement of P2X4 and P2Y12 receptors in ATP-induced microglial chemotaxis. *Glia.* 55: 604-616, 2007.

Oike M, Droogmans G, Ito Y. ATP release pathways in vascular endothelial cells. *Nippon Yakurigaku Zasshi*. 123: 403-411, 2004.

Osborne CK, Hobbs K, Trent JM. Biological differences among MCF-7 human breast cancer cell lines from different laboratories. *Breast Cancer Res. Treat.*, 9: 111-121, 1987.

Park S, Mazina O, Kitagawa A, Wong P, Matsumura F. TCDD causes suppression of growth and differentiation of MCF10A, human mammary epithelial cells by interfering with their insulin receptor signaling through c-Src kinase and ERK activation. *J Biochem Mol Toxicol*. 18: 322-331, 2004.

Pearson GW, Earnest S, Cobb MH. Cyclic AMP selectively uncouples MAP kinase cascades from activating signals. *Mol Cell Biol*. 26:3039-3047, 2006.

Potucek ID, Crain JM, Watters JJ. Purinergic receptors modulate MAP kinases and transcription factors that control microglial inflammatory gene expression. *Neurochem. Int*. 49:204-214, 2006.

Ralevic V, Burnstock G. Receptors for Purines and Pyrimidines. *Pharmacol Rev*. 50: 413- 492, 1998.

Raman M, Chen W, Cobb MH. Differential regulation and properties of MAPKs. *Oncogene*. 26: 3100-3112, 2007.

Ronda AC, Buitrago C, Colicheo A, de Boland AR, Roldán E, Boland R. Activation of MAPKs by 1 α ,25(OH) $_2$ -vitamin D $_3$ and 17 β -Estradiol in skeletal muscle cells leads to phosphorylation of Elk-1 and CREB transcription factors. *J Steroid Biochem Mol Biol*. 103: 462-466, 2007.

Roskoski R. Src protein-tyrosine kinase structure and regulation. *Biochem Biophys Res Commun*. 324: 1155-1164, 2004.

Roskoski R. Src kinase regulation by phosphorylation and dephosphorylation. *Biochem Biophys Res Commun*. 331: 1-14, 2005.

Rubin J, Rubin C, Jacobs CR. Molecular pathways mediating mechanical signaling in bone. *Gene*. 15: 1-16, 2006.

Sabapathy K, Hochedlinger K, Nam SY, Bauer A, Karin M, Wagner EF. Distinct roles for JNK1 and JNK2 in regulating JNK activity and c-Jun-dependent cell proliferation. *Mol Cell*. 15: 713-725, 2004.

Sadoshima J, Izumo S. Mechanotransduction in stretch-induced hypertrophy of cardiac myocytes. *J Recept Res*. 13: 777-794, 1993.

Salamanca DA, Khalil RA. Protein kinase C isoformes as specific targets for modulation of vascular smooth muscle function in hypertension. *Biochem Pharmacol.* 70: 1537-1547, 2005.

Salter MW, Hicks JL. ATP causes release of intracellular Ca²⁺ via the phospholipase C β /IP₃ pathway in astrocytes from the dorsal spinal cord. *J Neurosci.* 15: 2961-2967, 1995.

Sanchez-Esteban J, Cicchiello LA, Wang Y, Tsai SW, Williams LK, Torday JS, Rubin LP. Mechanical stretch promotes alveolar epithelial type II cell differentiation. *J Appl Physiol.* 91: 589-595, 2001.

Sanchez-Prieto R, Rojas JM, Taya Y, Gutkind JS. A role for the p38 mitogen-activated protein kinase pathway in the transcriptional activation of p53 on genotoxic stress by chemotherapeutic agents. *Cancer Res.* 60: 2464-2472, 2000.

Santillán G, Baldi C, Katz S, Vazquez G, Boland R. Evidence that TRPC3 is a molecular component of the 1 α , 25(OH)₂ D₃-activated capacitative calcium entry (CCE) in muscle and osteoblast cells. *J. Steroid Biochem. Mol. Biol.* 89/90: 291-295, 2004.

Schafer R, Sedehizade F, Welte T, Reiser G. ATP- and UTP-activated P₂Y receptors differently regulate proliferation of human lung epithelial tumor cells. *Am J Physiol Lung Cell Mol Physiol.* 285: L376-L385, 2003.

Scodelaro Bilbao P, Boland R, de Boland AR, Santillán G. ATP modulation of Mitogen Activated Protein Kinases and intracellular Ca²⁺ in breast cancer (MCF-7) cells. *Arch. Biochem. Biophys.* 466: 15-23, 2007.

Sellers LA, Simon J, Lundahl TS, Cousens DJ, Humphrey PPA, Barnard EA. Adenosine nucleotides acting at the human P₂Y₁ receptor stimulate mitogen-activated protein kinases and induce apoptosis. *J Biol Chem.* 276: 16379-16390, 2001.

Sharma RV, Chapleau MW, Hajduczuk G, Wachtel RE, Waite LJ, Bhalla RC, Abboud FM. Mechanical stimulation increases intracellular calcium concentration in nodose sensory neurons. *Neuroscience.* 66: 433-441, 1995.

Simon C, Goepfert H, Boyd D. Inhibition of the p38 mitogen-activated protein kinase by SB 203580 blocks PMA-induced Mr 92,000 type IV

- collagenase secretion and in vitro invasion. *Cancer Res.* 58: 1135-1139, 1998.
- Smeal T, Angel P, Meek J, Karin M. Different requirements for formation of Jun:Jun and Jun:Fos complexes. *Genes Develop.* 3: 2091-2100, 1989.
- Snedecor GW, Cochran WG. Statistical methods. Ames,IA: Iowa State University Press, 1967.
- Sonnenburg ED, Gao T, Newton AC. The Phosphoinositide-dependent Kinase, PDK-1, phosphorylates conventional Protein Kinase C isozymes by a mechanism that is independent of Phosphoinositide 3-Kinase. *J Biol Chem.* 276: 45289-45297, 2001.
- Syed NA, Horner KN, Misra V, Khandelwal RL. Different cellular localization, translocation, and insulin-induced phosphorylation of PKB α in HepG2 cells and hepatocytes. *J Cell Biochem.* 86: 118-127, 2002.
- Thevananther S, Sun H, Li D, Arjunan V, Awad SS, Wyllie S, Zimmerman TL, Goss JA, Karpen SJ. Extracellular ATP activates c-jun N-terminal kinase signaling and cell cycle progression in hepatocytes. *Hepatology.* 39: 393-402, 2004.
- Thiel G, Sarraj JA, Vinson C, Stefano L, Bach K. Role of basic region leucine zipper transcription factors cyclic AMP response element binding protein (CREB), CREB2, activating transcription factor 2 and CAAT/enhancer binding protein α in cyclic AMP response element mediated transcription. *J Neurochem.* 92: 321-336, 2005.
- Tinel H, Denker HW, Thie M. Calcium influx in human uterine epithelial RL95-2 cells triggers adhesiveness for trophoblast-like cells. Model studies on signalling events during embryo implantation. *Mol Hum Reprod.* 6: 1119-1130, 2000.
- Towbin H, Staehelin T, Gordon J. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc Natl Acad Sci.* 76: 4350-4354, 1979.
- Tracy S, van der Geer P, Hunter T. The receptor-like protein-tyrosine phosphatase, RPTP α , is phosphorylated by protein kinase C on two serines close to the inner face of the plasma membrane. *J Biol Chem.* 270: 10587-10594, 1995.
- Tu MT, Luo SF, Wang CC, Chien CS, Chiu CT, Lin CC, Yang CM. P2Y₂ receptor-mediated proliferation of C₆ glioma cells via activation of

- Ras/Raf/MEK/MAPK pathway. *Br. J. Pharmacol.* 129: 1481-1489, 2000.
- Turjanski AG, Vaqué JP, Gutkind JS. MAP kinases and the control of nuclear events. *Oncogene.* 26: 3240-3253, 2007.
- van Biesen T, Luttrell LM, Hawes BE, Lefkowitz RJ. Mitogenic signaling via G protein-coupled receptors. *Endocr. Rev.* 17: 698-714, 1996.
- Vanhaesebroeck B, Alessi DR. The PI3K-PDK1 connection: more than just a road to PKB. *Biochem J.* 346: 561-76, 2000.
- van Kolen K, Slegers H. Integration of P2Y receptor-activated signal transduction pathways in G protein-dependent signaling networks. *Purinergic Signal.* 2: 451-469, 2006.
- von Kugelgen I. Pharmacological profiles of cloned mammalian P2Y-receptor subtypes. *Pharmacol. & Ther.* 110: 415-432, 2006.
- Wagstaff SC, Bowler WB, Gallagher JA, Hipskind RA. Extracellular ATP activates multiple signaling pathways and potentiates growth factor-induced c-Fos gene expression in MCF-7 cells. *Carcinogenesis.* 21: 2175-2181, 2000.
- Wan M-X, Ren L-M, Shan B-E. Inhibitory effects of extracellular adenosine triphosphate on growth of esophageal carcinoma cells. *World J Gastroenterol.* 11: 5915-5919, 2005.
- White N, Burnstock G. P2 receptors and cancer. *TRENDS in Pharmacol Sci.* 27: 211-217, 2006.
- Wu J, Kamimura N, Takeo T, Suga S, Wakui M, Maruyama T, Mikoshiba K. 2-Aminoethoxydiphenyl borate modulates kinetics of intracellular Ca²⁺ signals mediated by inositol 1,4,5-trisphosphate-sensitive Ca²⁺ stores in single pancreatic acinar cells of mouse. *Mol Pharmacol.* 58: 1368-74, 2000.
- Wu M, Holowka D, Craighead HG, Baird B. Visualization of plasma membrane compartmentalization with patterned lipid bilayers. *PNAS.* 101: 13798-13803, 2004.
- Yang SH, Wu ZZ, Chien CM, Lo YH, Wu MJ, Chang LS, Lin SR. JNK and ERK mitogen-activated protein kinases mediate THDA-induced apoptosis in K562 cells. *Cell Biol Toxicol.* 24: 291-302, 2008.

- Yano S, Komine M, Fujimoto M, Okochi H, Tamaki K. Mechanical stretching in vitro regulates signal transduction pathways and cellular proliferation in human epidermal keratinocytes. *J Invest Dermatol.* 122: 783–790, 2004.
- Yates S, Rayner TE. Transcription factor activation in response to cutaneous injury: role of AP-1 in reepithelialization. *Wound Repair Regen.* 10: 5–15, 2002.
- Yin J, Xu K, Zhang J, Kumar A, Yu F-S. Wound-induced ATP release and EGF receptor activation in epithelial cells. *J Cell Sci.* 120: 815–825, 2007.
- Ying B, Fan H, Wen F, Xu D, Liu D, Yang D, Chen G, Dou L, Jiang F. Mechanical strain-induced c-fos expression in pulmonary epithelial cell line A549. *Biochem Biophys Res Commun.* 347: 369–372, 2006.
- Zarubin T, Han J. Activation and signaling of the p38 MAP kinase pathway. *Cell Res.* 15: 11–18, 2005.
- Zhang Y, Dong C. Regulatory mechanisms of mitogen-activated kinase signaling. *Cell Mol Life Sci.* 64: 2771–2789, 2007.
- Zhou H, Iwasaki H, Nakamura T, Nakamura K, Maruyama T, Hamano S, Ozaki S, Mizutani A, Mikoshiba K. 2-Aminoethyl diphenylborinate analogues: selective inhibition for store-operated Ca^{2+} entry. *Biochem Biophys Res Commun.* 352: 277–82, 2007.
- Zhu J, Blenis J, Yuan J. Activation of PI3K/Akt and MAPK pathways regulates Myc-mediated transcription by phosphorylating and promoting the degradation of Mad1. *Proc Natl Acad Sci.* 105: 6584–6589, 2008.