

## REFERENCIAS

- Abrams L, Politi LE, Adler R. Differential susceptibility of isolated mouse retinal neurons and photoreceptors to kainic acid toxicity. *In vitro* studies. *Invest Ophthalmol.Vis.Sci.* 1989; 30; 2300-2308.
- Acharya JK, Dasgupta U, Rawat SS, Yuan C, Sanxaridis PD, Yonamine I, Karim P, Nagashima K, Brodsky MH, Tsunoda S, Acharya U. Cell-nonautonomous function of ceramidase in photoreceptor homeostasis. *Neuron.* 2008;57(1):69-79.
- Acharya U, Mowen MB, Nagashima K, Acharya JK. Ceramidase expression facilitates membrane turnover and endocytosis of rhodopsin in photoreceptors. *Proc Natl Acad Sci U S A.* 2004;101(7):1922-6.
- Acharya U, Patel S, Koundakjian E, Nagashima K, Han X, Acharya JK. Modulating sphingolipid biosynthetic pathway rescues photoreceptor degeneration. *Science.* 2003; 299(5613):1740-3.
- Adler R, Canto-Soler MV. Molecular mechanisms of optic vesicle development: complexities, ambiguities and controversies. *Dev Biol.* 2007; 305(1):1-13.
- Adler R. Regulation of neurite growth in purified retina neuronal cultures: effects of PNPf, a substratum-bound, neurite-promoting factor. *J Neurosci Res.* 1982; 8(2-3):165-77.
- Akagawa K. Barnstable CJ. Identification and characterization of cell types in monolayer cultures of rat retina using monoclonal antibodies. *Brain Res.* 1986; 383: 110-120.
- Akbar M, Calderon F, Wen Z, Kim HY. Docosahexaenoic acid: a positive modulator of Akt signaling in neuronal survival. *Proc Natl Acad Sci U S A.* 2005; 102(31):10858-63.
- Akbar M, Kim HY. Protective effects of docosahexaenoic acid in staurosporine-induced apoptosis: involvement of phosphatidylinositol-3 kinase pathway. *J Neurochem.* 2002; 82(3):655-65.
- Anderson RE, Maude MB, McClellan M, Matthes MT, Yasumura D, LaVail MM. Low docosahexaenoic acid levels in rod outer segments of rats with P23H and S334ter rhodopsin mutations. *Mol Vis.* 2002; 8:351-8.
- Andrieu-Abadie N, Gouazé V, Salvayre R, Levade T. Ceramide in apoptosis signaling: relationship with oxidative stress. *Free Radic Biol Med.* 2001; 31(6):717-28.
- Annis MG, Soucie EL, Dlugosz PJ, Cruz-Aguado JA, Penn LZ, Leber B, Andrews DW. Bax forms multispinning monomers that oligomerize to permeabilize membranes during apoptosis. *EMBO J.* 2005; 24(12):2096-103.
- Antonetti DA, Barber AJ, Bronson SK, Freeman WM, Gardner TW, Jefferson LS, Kester M, Kimball SR, Krady JK, LaNoue KF, Norbury CC, Quinn PG, Sandrasegaran L, Simpson IA. Diabetic retinopathy: seeing beyond glucose-induced microvascular disease. *Diabetes.* 2006; 55(9):2401-11.
- Arab K, Rossary A, Flourié F, Tourneur Y, Steghens JP. Docosahexaenoic acid enhances the antioxidant response of human fibroblasts by upregulating gamma-glutamyl-cysteinyl ligase and glutathione reductase. *Br J Nutr.* 2006; 95(1):18-26.
- Archibald NK, Clarke MP, Mosimann UP, Burn DJ. The retina in Parkinson's disease. *Brain.* 2009; 132(Pt 5):1128-45.

- Ariga T, Jarvis WD, Yu RK. Role of sphingolipid-mediated cell death in neurodegenerative diseases. *J Lipid Res.* 1998; 39(1):1-16.
- Barber AJ. A new view of diabetic retinopathy: a neurodegenerative disease of the eye. *Prog Neuropsychopharmacol Biol Psychiatry.* 2003; 27(2):283-90.
- Barcellona ML, Cardiel G, Gratton E. Time-resolved fluorescence of DAPI in solution and bound to polydeoxynucleotides. *Biochem Biophys Res Commun.* 1990; 170(1):270-80.
- Barnstable CJ, Hofstein R, Akagawa K. A marker of early amacrine cell development in rat retina. *Brain Res.* 1985; 352: 286-290.
- Barnstable CJ. Immunological studies of the diversity and development of the mammalian visual system. *Immunol Rev.* 1987; 100:47-78.
- Baron W, Metz B, Bansal R, Hoekstra D, de Vries H. PDGF and FGF-2 signaling in oligodendrocyte progenitor cells: regulation of proliferation and differentiation by multiple intracellular signaling pathways. *Mol Cell Neurosci.* 2000; 15(3):314-29.
- Bassi R, Anelli V, Giussani P, Tettamanti G, Viani P, Riboni L. Sphingosine-1-phosphate is released by cerebellar astrocytes in response to bFGF and induces astrocyte proliferation through Gi-protein-coupled receptors. *Glia.* 2006; 53(6):621-30.
- Bazan NG, Scott BL, Reddy TS, Pelias MZ. Decreased content of docosahexaenoate and arachidonate in plasma phospholipids in Usher's syndrome. *Biochem Biophys Res Commun.* 1986; 141(2):600-4.
- Bazan NG. Cell survival matters: docosahexaenoic acid signaling, neuroprotection and photoreceptors. *Trends Neurosci.* 2006; 29(5):263-71.
- Bazzi MD, Nelsestuen GL. Mechanism of protein kinase C inhibition by sphingosine. *Biochem Biophys Res Commun.* 1987; 146(1):203-7.
- Ben-Yoseph O, Boxer PA, Ross BD. Assessment of the role of the glutathione and pentose phosphate pathways in the protection of primary cerebrocortical cultures from oxidative stress. *J Neurochem.* 1996; 66(6):2329-37.
- Betito S, Cuvillier O. Regulation by sphingosine 1-phosphate of Bax and Bad activities during apoptosis in a MEK-dependent manner. *Biochem Biophys Res Commun.* 2006; 340(4):1273-7.
- Bose R, Verheij M, Haimovitz-Friedman A, Scotto K, Fuks Z, Kolesnick R. Ceramide synthase mediates daunorubicin-induced apoptosis: an alternative mechanism for generating death signals. *Cell.* 1995; 82(3):405-14.
- Bottenstein JE, Sato GH. Growth of a rat neuroblastoma cell line in serum-free supplemented medium. *Proc Natl Acad Sci U S A.* 1979; 76(1):514-7.
- Braak M, Danneberg K, Lichte K, Liphardt K, Ktistakis NT, Pitson SM, Hla T, Jakobs KH, Meyer zu Heringdorf D. Galpha(q)-mediated plasma membrane translocation of sphingosine kinase-1 and cross-activation of S1P receptors. *Biochim Biophys Acta.* 2009; 1791(5):357-70.
- Bras J, Singleton A, Cookson MR, Hardy J. Emerging pathways in genetic Parkinson's disease: Potential role of ceramide metabolism in Lewy body disease. *FEBS J.* 2008; 275(23):5767-73.

- Bringmann A, Iandiev I, Pannicke T, Wurm A, Hollborn M, Wiedemann P, Osborne NN, Reichenbach A. Cellular signaling and factors involved in Müller cell gliosis: neuroprotective and detrimental effects. *Prog Retin Eye Res.* 2009; 28(6):423-51.
- Bringmann A, Reichenbach A. Role of Muller cells in retinal degenerations. *Front Biosci.* 2001; 6:E72-92.
- Brizuela L, Rábano M, Gangoiti P, Narbona N, Macarulla JM, Trueba M, Gómez-Muñoz A. Sphingosine-1-phosphate stimulates aldosterone secretion through a mechanism involving the PI3K/PKB and MEK/ERK 1/2 pathways. *J Lipid Res.* 2007; 48(10):2264-74.
- Bryan L, Kordula T, Spiegel S, Milstien S. Regulation and functions of sphingosine kinases in the brain. *Biochim Biophys Acta.* 2008; 1781(9):459-66.
- Buffo A, Rite I, Tripathi P, Lepier A, Colak D, Horn AP, Mori T, Götz M. Origin and progeny of reactive gliosis: A source of multipotent cells in the injured brain. *Proc Natl Acad Sci U S A.* 2008; 105(9):3581-6.
- Buniatian GH, Gebhardt R, Mecke D, Traub P, Wiesinger H. Common myofibroblastic features of newborn rat astrocytes and cirrhotic rat liver stellate cells in early cultures and *in vivo*. *Neurochem Int.* 1999; 35(4):317-27.
- Cao W, Li F, Steinberg RH, Lavail MM. Development of normal and injury-induced gene expression of aFGF, bFGF, CNTF, BDNF, GFAP and IGF-I in the rat retina. *Exp Eye Res.* 2001; 72(5):591-604.
- Castello PR, Drechsel DA, Day BJ, Patel M. Inhibition of mitochondrial hydrogen peroxide production by lipophilic metalloporphyrins. *J Pharmacol Exp Ther.* 2008; 324(3):970-6.
- Castello PR, Drechsel DA, Patel M. Mitochondria are a major source of paraquat-induced reactive oxygen species production in the brain. *J Biol Chem.* 2007; 282(19):14186-93.
- Catarzi S, Degl'Innocenti D, Iantomasi T, Favilli F, Vincenzini MT. The role of H<sub>2</sub>O<sub>2</sub> in the platelet-derived growth factor-induced transcription of the gamma-glutamylcysteine synthetase heavy subunit. *Cell Mol Life Sci.* 2002; 59(8):1388-94.
- Chang GQ, Hao Y, Wong F. Apoptosis: final common pathway of photoreceptor death in rd, rds, and rhodopsin mutant mice. *Neuron.* 1993; 11(4):595-605.
- Cho E, Hung S, Willett WC, Spiegelman D, Rimm EB, Seddon JM, Colditz GA, Hankinson SE. Prospective study of dietary fat and the risk of age-related macular degeneration. *Am J Clin Nutr.* 2001; 73(2):209-18.
- Choi MS, Anderson MA, Zhang Z, Zimonjic DB, Popescu N, Mukherjee AB. Neutral ceramidase gene: role in regulating ceramide-induced apoptosis. *Gene.* 2003; 315:113-22.
- Chucair AJ, Rotstein NP, Sangiovanni JP, During A, Chew EY, Politi LE. Lutein and zeaxanthin protect photoreceptors from apoptosis induced by oxidative stress: relation with docosahexaenoic acid. *Invest Ophthalmol Vis Sci.* 2007; 48(11):5168-77.
- Close JL, Reh TA. Regeneration: transdifferentiation and stem cells en Sernagor E, Eglén S, Harris B, Wong C. *Retinal Development* 307-317 pp. Ed. Cambridge University. UK. 2006.
- Coelho RP, Saini HS, Sato-Bigbee C. Sphingosine-1-phosphate and oligodendrocytes: From cell development to the treatment of multiple sclerosis. *Prostaglandins Other Lipid Mediat.* 2009.

- Cottet S, Schorderet DF. Mechanisms of apoptosis in retinitis pigmentosa. *Curr.Mol.Med.* 2009, 9; 375-383.
- Curcio CA, Sloan KR, Kalina RE, Hendrickson AE. Human photoreceptor topography. *J Comp Neurol.* 1990; 292(4):497-523.
- Cuvillier O, Edsall L, Spiegel S. Involvement of sphingosine in mitochondria-dependent Fas-induced apoptosis of type II Jurkat T cells. *J Biol Chem.* 2000; 275(21):15691-700.
- Cuvillier O, Levade T. Sphingosine 1-phosphate antagonizes apoptosis of human leukemia cells by inhibiting release of cytochrome c and Smac/DIABLO from mitochondria. *Blood.* 2001; 98(9): 2828-36.
- Cuvillier O, Nava VE, Murthy SK, Edsall LC, Levade T, Milstien S, Spiegel S. Sphingosine generation, cytochrome c release, and activation of caspase-7 in doxorubicin-induced apoptosis of MCF7 breast adenocarcinoma cells. *Cell Death Differ.* 2001; 8(2):162-71.
- Cuvillier O, Nava VE, Murthy SK, Edsall LC, Levade T, Milstien S, Spiegel S. Sphingosine generation, cytochrome c release, and activation of caspase-7 in doxorubicin-induced apoptosis of MCF7 breast adenocarcinoma cells. *Cell Death Differ.* 2001; 8(2):162-71.
- Cuvillier O, Pirianov G, Kleuser B, Vanek PG, Coso OA, Gutkind S, Spiegel S. Suppression of ceramide-mediated programmed cell death by sphingosine-1-phosphate. *Nature.* 1996; 381(6585):800-3.
- Cuvillier O. Sphingosine in apoptosis signaling. *Biochim Biophys Acta.* 2002; 1585(2-3):153-62.
- De La Paz M, Anderson RE. Region and age-dependent variation in susceptibility of the human retina to lipid peroxidation. *Invest Ophthalmol Vis Sci.* 1992; Vol 33, 3497-3499.
- Deigner HP, Haberkorn U, Kinscherf R. Apoptosis modulators in the therapy of neurodegenerative diseases. *Expert Opin Investig Drugs.* 2000; 9(4):747-64.
- Deretic D, Traverso V, Parkins N, Jackson F, Rodriguez de Turco EB, Ransom N. Phosphoinositides, ezrin/moesin, and rac1 regulate fusion of rhodopsin transport carriers in retinal photoreceptors. *Mol Biol Cell.* 2004; 15(1):359-70.
- Dev KK, Mullershausen F, Mattes H, Kuhn RR, Bilbe G, Hoyer D, Mir A. Brain sphingosine-1-phosphate receptors: implication for FTY720 in the treatment of multiple sclerosis. *Pharmacol Ther.* 2008; 117(1):77-93.
- Dyer MA, Cepko CL. Control of Müller glial cell proliferation and activation following retinal injury. *Nat Neurosci.* 2000; 3(9):873-80.
- Elrick MJ, Fluss S, Colombini M. Sphingosine, a product of ceramide hydrolysis, influences the formation of ceramide channels. *Biophys J.* 2006; 91(5):1749-56.
- Emerit J, Edeas M, Bricaire F. Neurodegenerative diseases and oxidative stress. *Biomed Pharmacother.* 2004; 58(1):39-46.
- Fain GL. Why photoreceptors die (and why they don't). *Bioessays.* 2006; 28(4):344-54.
- Fischer AJ, Omar G, Eubanks J, McGuire CR, Dierks BD, Reh TA. Different aspects of gliosis in retinal Muller glia can be induced by CNTF, insulin, and FGF2 in the absence of damage. *Mol Vis.* 2004; 10:973-86.

- Fischer AJ, Reh TA. Müller glia are a potential source of neural regeneration in the postnatal chicken retina. *Nat Neurosci.* 2001; 4(3):247-52.
- Fischer AJ, Reh TA. Potential of Müller glia to become neurogenic retinal progenitor cells. *Glia.* 2003; 43(1):70-6.
- Folch J, Lees M, Sloane Stanley GH. A simple method for the isolation and purification of total lipides from animal tissues. *J Biol Chem.* 1957; 226(1):497-509.
- García-Pacios M, Collado MI, Busto JV, Sot J, Alonso A, Arrondo JL, Goñi FM. Sphingosine-1-phosphate as an amphipathic metabolite: its properties in aqueous and membrane environments. *Biophys J.* 2009; 97(5):1398-407.
- García-Ruiz C, Colell A, Marí M, Morales A, Fernández-Checa JC. Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species. Role of mitochondrial glutathione. *J Biol Chem.* 1997; 272(17):11369-77.
- Gennero I, Fauvel J, Nieto M, Cariven C, Gaits F, Briand-Mésange F, Chap H, Salles JP. Apoptotic effect of sphingosine 1-phosphate and increased sphingosine 1-phosphate hydrolysis on mesangial cells cultured at low cell density. *J Biol Chem.* 2002; 277(15):12724-34.
- George-Weinstein M, Foster RF, Gerhart JV, Kaufman SJ. *In vitro* and *in vivo* expression of alpha 7 integrin and desmin define the primary and secondary myogenic lineages. *Dev Biol.* 1993; 156(1):209-29.
- German OL, Insua MF, Gentili C, Rotstein NP, Politi LE. Docosahexaenoic acid prevents apoptosis of retina photoreceptors by activating the ERK/MAPK pathway. *J Neurochem.* 2006; 98(5):1507-20.
- Gouazé V, Mirault ME, Carpentier S, Salvayre R, Levade T, Andrieu-Abadie N. Glutathione peroxidase-1 overexpression prevents ceramide production and partially inhibits apoptosis in doxorubicin-treated human breast carcinoma cells. *Mol Pharmacol.* 2001; 60(3):488-96.
- Guidry C. Isolation and characterization of porcine Müller cells. Myofibroblastic dedifferentiation in culture. *Invest Ophthalmol Vis Sci.* 1996; 37(5):740-52.
- Guo L, Duggan J, Cordeiro MF. Alzheimer's Disease and Retinal Neurodegeneration. *Curr Alzheimer Res.* 2009.
- Gupta N, Yücel YH. Glaucoma as a neurodegenerative disease. *Curr Opin Ophthalmol.* 2007; 18(2):110-4.
- Halliwell B. Oxidative stress and neurodegeneration: where are we now? *J Neurochem.* 2006; 97(6):1634-58.
- Hannun YA. The sphingomyelin cycle and the second messenger function of ceramide. *J Biol Chem.* 1994; 269(5):3125-8.
- Harada T, Harada C, Kohsaka S, Wada E, Yoshida K, Ohno S, Mamada H, Tanaka K, Parada LF, Wada K. Microglia-Müller glia cell interactions control neurotrophic factor production during light-induced retinal degeneration. *J Neurosci.* 2002; 22(21):9228-36.
- He X, Huang Y, Li B, Gong CX, Schuchman EH. Deregulation of sphingolipid metabolism in Alzheimer's disease. *Neurobiol Aging.* 2008.
- Hengartner MO. The biochemistry of apoptosis. *Nature.* 2000; 407(6805):770-6.

- Hitchcock P, Ochocinska M, Sieh A, Otteson D. Persistent and injury-induced neurogenesis in the vertebrate retina. *Prog Retin Eye Res.* 2004; 23(2):183-94.
- Hoffman DR, Birch DG. Docosahexaenoic acid in red blood cells of patients with X-linked retinitis pigmentosa. *Invest Ophthalmol Vis Sci.* 1995; 36(6):1009-18.
- Honjo M, Tanihara H, Kido N, Inatani M, Okazaki K, Honda Y. Expression of ciliary neurotrophic factor activated by retinal Müller cells in eyes with NMDA- and kainic acid-induced neuronal death. *Invest Ophthalmol Vis Sci.* 2000; 41(2):552-60.
- Hsieh HL, Sun CC, Wu CB, Wu CY, Tung WH, Wang HH, Yang CM. Sphingosine 1-phosphate induces EGFR expression via Akt/NF-kappaB and ERK/AP-1 pathways in rat vascular smooth muscle cells. *J Cell Biochem.* 2008; 103(6):1732-46.
- Huster D, Reichenbach A, Reichelt W. The glutathione content of retinal Müller (glial) cells: effect of pathological conditions. *Neurochem Int.* 2000; 36(4-5):461-9.
- Igarashi J, Erwin PA, Dantas AP, Chen H, Michel T. VEGF induces S1P1 receptors in endothelial cells: Implications for cross-talk between sphingolipid and growth factor receptors. *Proc Natl Acad Sci U S A.* 2003; 100(19):10664-9.
- Jarrett SG, Lin H, Godley BF, Boulton ME. Mitochondrial DNA damage and its potential role in retinal degeneration. *Prog Retin Eye Res.* 2008; 27(6):596-607.
- Jarvis WD, Kolesnick RN, Fornari FA, Traylor RS, Gewirtz DA, Grant S. Induction of apoptotic DNA damage and cell death by activation of the sphingomyelin pathway. *Proc Natl Acad Sci U S A.* 1994; 91(1):73-7.
- Ju WK, Lee MY, Hofmann HD, Kirsch M, Chun MH. Expression of CNTF in Müller cells of the rat retina after pressure-induced ischemia. *Neuroreport.* 1999; 10(2):419-22.
- Kågedal K, Zhao M, Svensson I, Brunk UT. Sphingosine-induced apoptosis is dependent on lysosomal proteases. *Biochem J.* 2001; 359(Pt 2):335-43.
- Kalén A, Borchardt RA, Bell RM. Elevated ceramide levels in GH4C1 cells treated with retinoic acid. *Biochim Biophys Acta.* 1992; 1125(1):90-6.
- Katsel P, Li C, Haroutunian V. Gene expression alterations in the sphingolipid metabolism pathways during progression of dementia and Alzheimer's disease: a shift toward ceramide accumulation at the earliest recognizable stages of Alzheimer's disease? *Neurochem Res.* 2007; 32(4-5):845-56.
- Kitatani K, Idkowiak-Baldys J, Hannun YA. The sphingolipid salvage pathway in ceramide metabolism and signaling. *Cell Signal.* 2008; 20(6):1010-8.
- Kolb H, Fernández E, Nelson R. *Webvision: The organization of the retina and visual system.* University of Utah, 2002.
- Kolb H, Linberg KA, Fisher SK. Neurons of the human retina: a Golgi study. *J Comp Neurol.* 1992; 318(2):147-87.
- Kolesnick RN, Haimovitz-Friedman A, Fuks Z. The sphingomyelin signal transduction pathway mediates apoptosis for tumor necrosis factor, Fas, and ionizing radiation. *Biochem Cell Biol.* 1994; 72(11-12):471-4.
- Komatsu W, Ishihara K, Murata M, Saito H, Shinohara K. Docosahexaenoic acid suppresses nitric oxide production and inducible nitric oxide synthase expression in interferon-gamma

- plus lipopolysaccharide-stimulated murine macrophages by inhibiting the oxidative stress. *Free Radic Biol Med.* 2003; 34(8):1006-16.
- Kubista M, Akerman B, Nordén B. Characterization of interaction between DNA and 4',6-diamidino-2-phenylindole by optical spectroscopy. *Biochemistry.* 1987; 26(14):4545-53.
- Kunchithapautham K, Rohrer B. Apoptosis and autophagy in photoreceptors exposed to oxidative stress. *Autophagy.* 2007; 3(5):433-41.
- La Vail MM. Survival of some photoreceptor cells in albino rats following long-term exposure to continuous light. *Invest Ophthalmol Vis Sci Vol.* 1976; 15, 64-70.
- Lauritzen I, Blondeau N, Heurteaux C, Widmann C, Romey G, Lazdunski M. Polyunsaturated fatty acids are potent neuroprotectors. *EMBO J.* 2000; 19(8):1784-93.
- Lebman DA, Spiegel S. Cross-talk at the crossroads of sphingosine-1-phosphate, growth factors, and cytokine signaling. *J Lipid Res.* 2008; 49(7):1388-94.
- Levin DE. Cell wall integrity signaling in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev.* 2005; 69(2):262-91.
- Limaye V. The role of sphingosine kinase and sphingosine-1-phosphate in the regulation of endothelial cell biology. *Endothelium.* 2008; 15(3):101-12.
- Lindskog M, Gleissman H, Ponthan F, Castro J, Kogner P, Johnsen JI. Neuroblastoma cell death in response to docosahexaenoic acid: sensitization to chemotherapy and arsenic-induced oxidative stress. *Int J Cancer.* 2006; 118(10):2584-93.
- Ma Y, Pitson S, Hercus T, Murphy J, Lopez A, Woodcock J. Sphingosine activates protein kinase A type II by a novel cAMP-independent mechanism. *J Biol Chem.* 2005; 280(28):26011-7.
- Maceyka M, Milstien S, Spiegel S. Shooting the messenger: oxidative stress regulates sphingosine-1-phosphate. *Circ Res.* 2007; 100(1):7-9.
- Maceyka M, Milstien S, Spiegel S. Sphingosine-1-phosphate: the Swiss army knife of sphingolipid signaling. *J Lipid Res.* 2009; 50 Suppl:S272-6.
- Maceyka M, Sankala H, Hait NC, Le Stunff H, Liu H, Toman R, Collier C, Zhang M, Satin LS, Merrill AH Jr, Milstien S, Spiegel S. SphK1 and SphK2, sphingosine kinase isoenzymes with opposing functions in sphingolipid metabolism. *J Biol Chem.* 2005; 280(44):37118-29.
- MacNeil MA, Masland RH. Extreme diversity among amacrine cells: implications for function. *Neuron.* 1998; 20(5):971-82.
- Mandal MA, Tran J, Zheng L, Brush RS. Sphingolipid Signaling in Retinal Cell Death. ARVO, Annual Meeting, 2009.
- Mann I. The development of the human eye. New York: Grune and Stratton; 1964.
- Masland RH. Neuronal diversity in the retina. *Curr Opin Neurobiol.* 2001; 11(4):431-6.
- McGille GS, Guidry C, Dacheux RF. Antigenic changes of rabbit retinal Müller cells in culture. *Invest Ophthalmol Vis Sci.* 1998; 39(8):1453-61.
- Means CK, Xiao CY, Li Z, Zhang T, Omens JH, Ishii I, Chun J, Brown JH. Sphingosine 1-phosphate S1P2 and S1P3 receptor-mediated Akt activation protects against *in vivo*

- myocardial ischemia-reperfusion injury. *Am J Physiol Heart Circ Physiol.* 2007; 292(6):H2944-51.
- Merrill AH Jr, Stokes TH, Momin A, Park H, Portz BJ, Kelly S, Wang E, Sullards MC, Wang MD. Sphingolipidomics: a valuable tool for understanding the roles of sphingolipids in biology and disease. *J Lipid Res.* 2009; 50 Suppl:S97-102.
- Merrill AH Jr. De novo sphingolipid biosynthesis: a necessary, but dangerous, pathway. *J Biol Chem.* 2002; 277(29):25843-6.
- Meyer zu Heringdorf D, Lass H, Kuchar I, Lipinski M, Alemany R, Rümenapp U, Jakobs KH. Stimulation of intracellular sphingosine-1-phosphate production by G-protein-coupled sphingosine-1-phosphate receptors. *Eur J Pharmacol.* 2001; 414(2-3):145-54.
- Milstien S, Gude D, Spiegel S. Sphingosine 1-phosphate in neural signalling and function. *Acta Paediatr Suppl.* 2007; 96(455):40-3.
- Miranda GE, Abraham CE, Politi LE, Rotstein NP. Sphingosine-1-phosphate is a key regulator of proliferation and differentiation in retina photoreceptors. *Invest Ophthalmol Vis Sci.* 2009; 50(9):4416-28.
- Mori TA, Woodman RJ, Burke V, Puddey IB, Croft KD, Beilin LJ. Effect of eicosapentaenoic acid and docosahexaenoic acid on oxidative stress and inflammatory markers in treated-hypertensive type 2 diabetic subjects. *Free Radic Biol Med.* 2003; 35(7):772-81.
- Moshiri A, Close J, Reh TA. Retinal stem cells and regeneration. *Int J Dev Biol.* 2004; 48(8-9):1003-14.
- Narayan S, Head SR, Gilmartin TJ, Dean B, Thomas EA. Evidence for disruption of sphingolipid metabolism in schizophrenia. *J Neurosci Res.* 2009; 87(1):278-88.
- Newman E, Reichenbach A. The Müller cell: a functional element of the retina. *Trends Neurosci.* 1996; 19(8):307-12.
- Ohta H, Sweeney EA, Masamune A, Yatomi Y, Hakomori S, Igarashi Y. Induction of apoptosis by sphingosine in human leukemic HL-60 cells: a possible endogenous modulator of apoptotic DNA fragmentation occurring during phorbol ester-induced differentiation. *Cancer Res.* 1995; 55(3):691-7.
- Olivera A, Spiegel S. Sphingosine-1-phosphate as second messenger in cell proliferation induced by PDGF and FCS mitogens. *Nature.* 1993;365(6446):557-60.
- Ooto S, Akagi T, Kageyama R, Akita J, Mandai M, Honda Y, Takahashi M. Potential for neural regeneration after neurotoxic injury in the adult mammalian retina. *Proc Natl Acad Sci U S A.* 2004; 101(37):13654-9.
- Ott M, Gogvadze V, Orrenius S, Zhivotovsky B. Mitochondria, oxidative stress and cell death. *Apoptosis.* 2007; 12:913-922.
- Oyama O, Sugimoto N, Qi X, Takuwa N, Mizugishi K, Koizumi J, Takuwa Y. The lysophospholipid mediator sphingosine-1-phosphate promotes angiogenesis *in vivo* in ischaemic hindlimbs of mice. *Cardiovasc Res.* 2008; 78(2):301-7.
- Park SH, Park JW, Park SJ, Kim KY, Chung JW, Chun MH, Oh SJ. Apoptotic death of photoreceptors in the streptozotocin-induced diabetic rat retina. *Diabetologia.* 2003; 46(9):1260-8.



- Pchejetski D, Kunduzova O, Dayon A, Calise D, Seguelas MH, Leducq N, Seif I, Parini A, Cu villier O. Oxidative stress-dependent sphingosine kinase-1 inhibition mediates monoamine oxidase A-associated cardiac cell apoptosis. *Circ Res.* 2007; 100(1):41-9.
- Pébay A, Toutant M, Prémont J, Calvo CF, Venance L, Cordier J, Glowinski J, Tencé M. Sphingosine-1-phosphate induces proliferation of astrocytes: regulation by intracellular signalling cascades. *Eur J Neurosci.* 2001; 13(12):2067-76.
- Polit L, Rotstein N, Carri N. Effects of docosahexaenoic acid on retinal development: cellular and molecular aspects. *Lipids.* 2001a; 36(9):927-35.
- Politi LE, Lehar M, Adler R. Development of neonatal mouse retinal neurons and photoreceptors in low density cell culture. *Invest Ophthalmol Vis Sci.* 1988; 29(4):534-43.
- Politi LE, Rotstein NP, Carri NG. Effect of GDNF on neuroblast proliferation and photoreceptor survival: additive protection with docosahexaenoic acid. *Invest Ophthalmol Vis Sci.* 2001b; 42(12):3008-15.
- Portera-Cailliau C, Sung CH, Nathans J, Adler R. Apoptotic photoreceptor cell death in mouse models of retinitis pigmentosa. *Proc Natl Acad Sci U S A.* 1994; 91(3):974-8.
- Pyne NJ, Long JS, Lee SC, Loveridge C, Gillies L, Pyne S. New Aspects of Sphingosine 1-phosphate Signaling in Mammalian Cells. *Adv Enzyme Regul.* 2009. En prensa.
- Quintyn JC, Ranty ML, Mathis A. Photoreceptor apoptosis in human retinal detachment. *Am J Ophthalmol.* 2006; 141(1):227.
- Raisova M, Goltz G, Bektas M, Bielawska A, Riebeling C, Hossini AM, Eberle J, Hannun YA, Orfanos CE, Geilen CC. Bcl-2 overexpression prevents apoptosis induced by ceramidase inhibitors in malignant melanoma and HaCaT keratinocytes. *FEBS Lett.* 2002; 516(1-3):47-52.
- Ranty ML, Carpentier S, Cournot M, Rico-Lattes I, Malecaze F, Levade T, Delisle MB, Quintyn JC. Ceramide production associated with retinal apoptosis after retinal detachment. *Graefes Arch Clin Exp Ophthalmol.* 2009; 247(2):215-24.
- Ravichandran KS, Lorenz U. Engulfment of apoptotic cells: signals for a good meal. *Nat Rev Immunol.* 2007; 7(12):964-74.
- Raymond MN, Bole-Feysot C, Banno Y, Tanfin Z, Robin P. Endothelin-1 inhibits apoptosis through a sphingosine kinase 1-dependent mechanism in uterine leiomyoma ELT3 cells. *Endocrinology.* 2006; 147(12):5873-82.
- Rius RA, Edsall LC, Spiegel S. Activation of sphingosine kinase in pheochromocytoma PC12 neuronal cells in response to trophic factors. *FEBS Lett.* 1997; 417(2):173-6.
- Rosen H, Goetzl EJ. Sphingosine 1-phosphate and its receptors: an autocrine and paracrine network. *Nat Rev Immunol.* 2005; 5(7):560-70.
- Rosen H, Gonzalez-Cabrera PJ, Sanna MG, Brown S. Sphingosine 1-phosphate receptor signaling. *Annu Rev Biochem.* 2009; 78:743-68.
- Rotstein NP, Aveldaño MI, Barrantes FJ, Politi LE. Docosahexaenoic acid is required for the survival of rat retinal photoreceptors *in vitro*. *J Neurochem.* 1996; 66(5):1851-9.

- Rotstein NP, Aveldaño MI, Barrantes FJ, Roccamo AM, Politi LE. Apoptosis of retinal photoreceptors during development *in vitro*: protective effect of docosahexaenoic acid. *J Neurochem.* 1997; 69(2):504-13.
- Rotstein NP, Politi LE, Aveldaño MI. Docosahexaenoic acid promotes differentiation of developing photoreceptors in culture. *Invest Ophthalmol.Vis.Sci.* 1998. 39; 2750-2758.
- Rotstein NP, Politi LE, German OL, Girotti R. Protective effect of docosahexaenoic acid on oxidative stress-induced apoptosis of retina photoreceptors. *Invest Ophthalmol Vis Sci.* 2003; 44(5):2252-9.
- Saini HS, Coelho RP, Goparaju SK, Jolly PS, Maceyka M, Spiegel S, Sato-Bigbee C. Novel role of sphingosine kinase 1 as a mediator of neurotrophin-3 action in oligodendrocyte progenitors. *J Neurochem.* 2005; 95(5):1298-310.
- Sakakura C, Sweeney EA, Shirahama T, Hagiwara A, Yamaguchi T, Takahashi T, Hakomori S, Igarashi Y. Selectivity of sphingosine-induced apoptosis. Lack of activity of DL-erythro-dihydrosphingosine. *Biochem Biophys Res Commun.* 1998; 246(3):827-30.
- Sanchez T, Hla T. Structural and functional characteristics of S1P receptors. *J Cell Biochem.* 2004; 92(5):913-22.
- Sanvicens N, Cotter TG. Ceramide is the key mediator of oxidative stress-induced apoptosis in retinal photoreceptor cells. *J Neurochem.* 2006; 98(5):1432-44.
- Sasaki H, Arai H, Cocco MJ, White SH. pH dependence of sphingosine aggregation. *Biophys J.* 2009; 96(7):2727-33.
- Sato K, Ishikawa K, Ui M, Okajima F. Sphingosine 1-phosphate induces expression of early growth response-1 and fibroblast growth factor-2 through mechanism involving extracellular signal-regulated kinase in astroglial cells. *Brain Res Mol Brain Res.* 1999; 74(1-2):182-9.
- Shimazawa M, Nakajima Y, Mashima Y, Hara H. Docosahexaenoic acid (DHA) has neuroprotective effects against oxidative stress in retinal ganglion cells. *Brain Res.* 2009; 1251:269-75.
- Siskind LJ, Fluss S, Bui M, Colombini M. Sphingosine forms channels in membranes that differ greatly from those formed by ceramide. *J Bioenerg Biomembr.* 2005; 37(4):227-36.
- Skoura A, Sanchez T, Claffey K, Mandala SM, Proia RL, Hla T. Essential role of sphingosine 1-phosphate receptor 2 in pathological angiogenesis of the mouse retina. *J Clin Invest.* 2007; 117(9):2506-16.
- Soldi R, Mandinova A, Venkataraman K, Hla T, Vadas M, Pitson S, Duarte M, Graziani I, Kolev V, Kacer D, Kirov A, Maciag T, Prudovsky I. Sphingosine kinase 1 is a critical component of the copper-dependent FGF1 export pathway. *Exp Cell Res.* 2007; 313(15):3308-18.
- Sorensen G, Medina S, Parchaliuk D, Phillipson C, Robertson C, Booth SA. Comprehensive transcriptional profiling of prion infection in mouse models reveals networks of responsive genes. *BMC Genomics.* 2008; 9:114.
- Spiegel S, Milstien S. Exogenous and intracellularly generated sphingosine 1-phosphate can regulate cellular processes by divergent pathways. *Biochem Soc Trans.* 2003; 31:1216-9.
- Spiegel S, Milstien S. Sphingosine 1-phosphate, a key cell signaling molecule. *J Biol Chem.* 2002; 277(29):25851-4.

- Spiegel S, Milstien S. Sphingosine-1-phosphate: an enigmatic signalling lipid. *Nat Rev Mol Cell Biol.* 2003a; 4(5):397-407.
- Suzuki E, Handa K, Toledo MS, Hakomori S. Sphingosine-dependent apoptosis: a unified concept based on multiple mechanisms operating in concert. *Proc Natl Acad Sci U S A.* 2004; 101 (41):14788-93.
- Swaney JS, Moreno KM, Gentile AM, Sabbadini RA, Stoller GL. Sphingosine-1-phosphate (S1P) is a novel fibrotic mediator in the eye. *Exp Eye Res.* 2008; 87(4):367-75.
- Sweeney EA, Inokuchi J, Igarashi Y. Inhibition of sphingolipid induced apoptosis by caspase inhibitors indicates that sphingosine acts in an earlier part of the apoptotic pathway than ceramide. *FEBS Lett.* 1998; 425(1):61-5.
- Taha TA, Argraves KM, Obeid LM. Sphingosine-1-phosphate receptors: receptor specificity versus functional redundancy. *Biochim Biophys Acta.* 2004; 1682(1-3):48-55.
- Taha TA, Mullen TD, Obeid LM. A house divided: ceramide, sphingosine, and sphingosine-1-phosphate in programmed cell death. *Biochim Biophys Acta.* 2006; 1758(12):2027-36.
- Theilmeyer G, Schmidt C, Herrmann J, Keul P, Schäfers M, Herrgott I, Mersmann J, Larmann J, Hermann S, Stypmann J, Schober O, Hildebrand R, Schulz R, Heusch G, Haude M, von Wnuck Lipinski K, Herzog C, Schmitz M, Erbel R, Chun J, Levkau B. High-density lipoproteins and their constituent, sphingosine-1-phosphate, directly protect the heart against ischemia/reperfusion injury *in vivo* via the S1P3 lysophospholipid receptor. *Circulation.* 2006; 114(13):1403-9.
- Vaney DI. The mosaic of amacrine cells in the mammalian retina. *Progress in retinal research.* 1990, 9, 49-100.
- Venkataraman K, Thangada S, Michaud J, Oo ML, Ai Y, Lee YM, Wu M, Parikh NS, Khan F, Proia RL, Hla T. Extracellular export of sphingosine kinase-1a contributes to the vascular S1P gradient. *Biochem J.* 2006 Aug 1; 397(3):461-71.
- Vercesi AE, Kowaltowski AJ, Oliveira HC, Castilho RF. Mitochondrial Ca<sup>2+</sup> transport, permeability transition and oxidative stress in cell death: implications in cardiotoxicity, neurodegeneration and dyslipidemias. *Front Biosci.* 2006; 11:2554-64.
- Wamhoff BR, Lynch KR, Macdonald TL, Owens GK. Sphingosine-1-phosphate receptor subtypes differentially regulate smooth muscle cell phenotype. *Arterioscler Thromb Vasc Biol.* 2008; 28(8):1454-61.
- Wan J, Zheng H, Xiao HL, She ZJ, Zhou GM. Sonic hedgehog promotes stem-cell potential of Müller glia in the mammalian retina. *Biochem Biophys Res Commun.* 2007; 363(2):347-54.
- Wang W, Graeler MH, Goetzl EJ. Type 4 sphingosine 1-phosphate G protein-coupled receptor (S1P4) transduces S1P effects on T cell proliferation and cytokine secretion without signaling migration. *FASEB J.* 2005; 19(12):1731-3.
- Wangsa-Wirawan ND, Linsenmeier RA. Retinal oxygen: fundamental and clinical aspects. *Arch Ophthalmol.* 2003; 121(4):547-57.
- Watabe S, Hiroi T, Yamamoto Y, Fujioka Y, Hasegawa H, Yago N, Takahashi SY. SP-22 is a thioredoxin-dependent peroxide reductase in mitochondria. *Eur J Biochem.* 1997; 249(1):52-60.

Wenzel A, Grimm C, Samardzija M, Remé CE. Molecular mechanisms of light-induced photoreceptor apoptosis and neuroprotection for retinal degeneration. *Prog Retin Eye Res.* 2005; 24(2):275-306.

Wiesner DA, Dawson G. Staurosporine induces programmed cell death in embryonic neurons and activation of the ceramide pathway. *J Neurochem.* 1996; 66(4):1418-25.

Woodcock J. Sphingosine and ceramide signalling in apoptosis. *IUBMB Life.* 2006; 58(8):462-6.

Woodcock JM, Murphy J, Stomski FC, Berndt MC, Lopez AF. The dimeric versus monomeric status of 14-3-3zeta is controlled by phosphorylation of Ser58 at the dimer interface. *J Biol Chem.* 2003; 278(38):36323-7.

Wu J, Bohanan CS, Neumann JC, Lingrel JB. KLF2 transcription factor modulates blood vessel maturation through smooth muscle cell migration. *J Biol Chem.* 2008; 283(7):3942-50.

Xia P, Wang L, Gamble JR, Vadas MA. Activation of sphingosine kinase by tumor necrosis factor-alpha inhibits apoptosis in human endothelial cells. *J Biol Chem.* 1999; 274(48):34499-505.

Zadro-Lamoureux LA, Zacks DN, Baker AN, Zheng QD, Hauswirth WW, Tsilfidis C. XIAP effects on retinal detachment-induced photoreceptor apoptosis. *Invest Ophthalmol Vis Sci.* 2009; 50(3):1448-53.

Zhu D, Sreekumar PG, Hinton DR, Kannan R. Expression and regulation of enzymes in the ceramide metabolic pathway in human retinal pigment epithelial cells and their relevance to retinal degeneration. *Vision Res.* 2009. En prensa.

**PUBLICACIONES ORIGINADAS DURANTE ESTA TESIS:**

- German OL, Miranda GE, Abrahan CE, Rotstein NP. Ceramide is a mediator of apoptosis in retina photoreceptors. *Invest Ophthalmol Vis Sci.* 2006; 47(4):1658-68.
- Abrahan CE, Insua MF, Politi LE, German OL, Rotstein NP. Oxidative stress promotes proliferation and dedifferentiation of retina glial cells *in vitro*. *J Neurosci Res.* 2009; 87(4):964-77.
- Miranda GE, Abrahan CE, Politi LE, Rotstein NP. Sphingosine-1-phosphate is a key regulator of proliferation and differentiation in retina photoreceptors. *Invest Ophthalmol Vis Sci.* 2009; 50(9):4416-28.
- Abrahan C, Miranda G, Agnolazza D, Politi L, Rotstein N. Synthesis of sphingosine is required for oxidative stress-induced apoptosis of photoreceptors. *Invest Ophthalmol Vis Sci.* 2009 Sep 24.