

Bibliografía

- Abrahams M V. (1986) Patch choice under perceptual constraints: a cause for departures from an ideal free distribution. *Behav Ecol Sociobiol* 19, 409-415.
- Akaike H. (1973) Information theory as an extension of the maximum likelihood principle. En: B.N. Petrov and F. Csaki (Eds), 2nd International Symposium on Information Theory. Akademiai Kiado, Budapest, p 267-281.
- Allee W C, Emerson A E, Park O, Park T y Schmidt K P. (1949) *Principles of Animal Ecology.*, 837 pp. W B Saunders, Philadelphia.
- Asher M, Spinelli de Oliveira E y Sachser N. (2004) Social system and spatial organization of wild guinea pigs (*Cavia aperea*) in a natural population. *Journal of Mammalogy* 85(4), 788-796.
- Baddeley A y Gill R D. (1997) Kaplan-Meier estimators of distance distributions for spatial point processes. *The Annals of Statistics* 25(1), 263-292.
- Baddeley A y Turner R. (2005) spatstat: an R package for analyzing spatial point patterns. *Journal of Statistical Software* 12(6), 1-42. ISSN: 1548-7660. URL: www.iostatsoft.org.
- Baldi R. (2007) Breeding success of the endemic mara *Dolichotis patagonum* in relation to habitat selection: Conservation implications. *Journal of Arid Environments* 68, 9-19.
- Barot S, Gignoux J y Menaut J C. (1999) Demography of a savanna palm tree: predictions from comprehensive spatial pattern analyses. *Ecology* 80(6), 1987-2005.
- Barros V. (1983) Atlas del potencial eólico de la Patagonia. Contribución N°. 69. Centro Nacional Patagónico, Puerto Madryn, Chubut, Argentina, 80 pp.
- Bates D, Maechler M y Bolker B. (2011) lme4: Linear mixed-effects models using Eigen and syntax. R package version 0.999375-39. <http://CRAN.R-project.org/package=lme4>.
- Battin J. (2004) When good animals love bad habitats: ecological traps and the conservation of animal populations. *Conservation Biology* . 18(6), 1482-1491.
- Beckmann J P y Berger J. (2003) Using black bears to test ideal-free distribution models experimentally. *Journal of Mammalogy* 84(2), 594-606.
- Beeskov A M, Elissalde N O y Rostagno C M. (1995) Ecosystem change associated with grazing intensity on the Punta Ninfa rangelands of Patagonia, Argentina. *Journal of Range Management* 48, 517-522.
- Beltramone C. (1981) Rasgos fisiográficos de Península Valdés. Contribución 46. CONICET. Centro Nacional Patagónico, Puerto Madryn, 18 pp.
- Ben-Shahar R y Skinner J D. (1988) Habitat preferences of african ungulates derived by uni- and multivariate analyses. *Ecology* 69(5), 1479-1485.
- Bertiller M B, Beeskov A M y Irrisari O. (1981) Caracteres fisonómicos y florísticos de la vegetación del Chubut. 2. La Península de Valdés y el Istmo Ameghino. Contribución 41. CONICET. Centro Nacional Patagónico, Puerto Madryn, 20 pp.
- Betts M G, Rodenhouse N L, Sillett T S, Doran P J y Holmes R T. (2008) Dynamic occupancy models reveal within-breeding season movement up a habitat quality gradient by a migratory songbird. *Ecography* 31, 592-600.
- Bilenca D N, Cittadino E A y Kravetz F O. (1995) Influencia de la actividad de *Cavia aperea* sobre la estructura del habitat y la distribución de *Akodon azarae* y *Oryzomys flavescens* (Rodentia: Caviidae, Muridae) en bordes de cultivos de la región pampeana (Argentina). *Iheringia, Série Zoologia, Porto Alegre* 79, 67-75.
- Bisigato A J y Bertiller M B. (1997) Grazing effects on patchy dryland vegetation in northern Patagonia. *Journal of Arid Environments* 36, 639-653.

- Bisigato A J, Bertiller M B, Ares J O y Pazos G E. (2005) Effect of grazing on plant patterns in arid ecosystems of Patagonian Monte. *Ecography* 28, 561-572.
- Bolker B y Pacala S W. (1997) Using moment equations to understand stochastically driven spatial pattern formation in ecological systems. *Theoretical Population Biology* 52, 179-197.
- Booth D T, Cox S E y Berryman R D. (2006) Point sampling digital imagery using 'SamplePoint'. *Environmental Monitoring and Assessment* 123, 97-108.
- Börger L, Dalziel B D y Fryxell J M. (2008) Are there general mechanisms of animal home range behaviour? A review and prospects for future research. *Ecology letters* 11(6), 637-650.
- Bowers M A y Barrett G W. (1999) Synthesis: a review of the science and prescriptions for the future. En: G.W. Barrett and J.D. Peles (Eds), *Landscape ecology of small mammals*, pp. 313-333. Springer-Verlag, New York.
- Bowman J, Forbes G J y Dilworth T G. (2001) The spatial component of variation in smallmammal abundance measured at three scales. *Canadian Journal of Zoology* 79, 137-144.
- Boyce M S, Mao J S, Merrill E H y Fortin D. (2003) Scale and heterogeneity in habitat selection by elk in Yellowstone National Park. *Ecoscience* 10(4), 421-431.
- Branch L C. (1993a) Intergroup and intragroup spacing in the plains vizcacha, *Lagostomus maximus*. *Journal of Mammalogy* 74(4), 890-900.
- Branch L C. (1993b) Social organization and mating system of the plains viscacha (*Lagostomus maximus*). *Journal of Zoology of London* 229, 473-491.
- Branch L C, Villarreal D y Fowler G S. (1993) Recruitment, dispersal, and group fusion in a declining population of the plains vizcacha (*Lagostomus maximus*; chinchillidae). *Journal of Mammalogy* 74(1), 9-20.
- Brown J. (1984) On the relationship between abundance and distribution of species. *The American Naturalist* 124(2), 255-279.
- Brown J H, Mehlman D W y Stevens G C. (1995) Spatial Variation in Abundance. *Ecology* 76, 2028-2043.
- Brown J H, Mehlman D W y Stevens G C. (2007) Spatial Variation in Abundance Emphasizing new ideas , to stimulate research in ecology. *Ecology* 76(7), 2028-2043.
- Bucher E. (1987) Herbivory in arid and semi-arid regions of Argentina. *Revista Chilena de Historia Natural* 60, 265-273.
- Campos C M, Tognelli M F y Ojeda R A. (2001) *Dolichotis Patagonum*. *Mammalian Species* 652, 1-5.
- Cañadas A, Sagarminaga R, De Stephanis R, Urquiola E y Hammond P S. (2005) Habitat preference modelling as a conservation tool: proposals for marine protected areas for cetaceans in southern Spanish waters. *Aquatic Conservation: Marine and Freshwater Ecosystems* 15, 495-521.
- Cheli G H. (2009) Efectos del disturbio por pastoreo ovino sobre la comunidad de artrópodos epígeos en Península Valdés (Chubut, Argentina). Doctoral, UNIVERSIDAD NACIONAL DEL COMAHUE.
- Clark P J y Evans F C. (1954) Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology* 35(4), 445-453.
- Condit R, Ashton P S, Baker P, Bunyavejchewin S, Gunatilleke S, Gunatilleke N, Hubbell S P, Foster R B, Itoh A, LaFrankie J V, Seng Lee H, Losos E, Manokaran N, Sukumar R y Yamakura T. (2000) Spatial patterns in the distribution of tropical tree species. *Science* 288(5470), 1414-1418.
- Conway C J, Garcia V, Smith M D, Ellis L A y Whitney J L. (2006) Comparative demography of burrowing owls in agricultural and urban landscapes in southeastern Washington. *Journal of Field Ornithology* 77, 280-290.
- Coronato F. (1993) Wind chill factor applied to Patagonian climatology. *International Journal of Biometeorological* 37(1), 1-6.

- Courchamp E, Clutton-Brock T y Grenfell B. (1999) Inverse density dependence and the Allee effect. *Trends in Ecology & Evolution* 14, 405-410.
- Dale M R T. (1995) Spatial pattern in communities of crustose saxicolous lichens. *Lichenologist* 27(6), 495-503.
- Danchin E y Wagner R H. (1997) The evolution of coloniality: the emergence of new perspectives. *TREE* 12(9), 342-347.
- Davis J M y Stamps J A. (2004) The effect of natal experience on habitat preferences. *TRENDS in Ecology and Evolution* 19(8), 411-416.
- Diggle P J. (1979) On Parameter Estimation and Goodness-of-Fit Testing for Spatial Point Patterns. *Biometrics* 35(1), 87-101.
- Djawdan M y Garland T. (1988) Maximal running speeds of bipedal and quadrupedal rodents. *Journal of Mammalogy* 69(4), 765-772.
- Donahue M J. (2006) Allee effects and conspecific cueing jointly lead to conspecific attraction. *Oecologia* 149, 33-43.
- Doncaster C P. (2000) Extension of ideal free resource use to breeding populations and metapopulations. *Oikos* 89, 24-36.
- Dunning J B, Danielson B J y Pulliam H R. (1992) Ecological processes that affect populations in complex landscapes. *Oikos* 65, 169-175.
- Ebensperger L A y Hayes L D. (2008) On the dynamics of rodent social groups. *Behavioural Processes* 79, 85-92.
- Ebensperger L A, Taraborelli P, Giannoni S M, Hurtado M J, León C y Bozinovic F. (2006) Nest and space use in a highland population of the southern mountain cavy (*Microcavia australis*). *Journal of Mammalogy* 87(5), 834-840.
- Elisalde N y Miravalles H. (1983) Evaluación de los campos de pastoreo de Península Valdés. Centro Nacional Patagónico. Contribución No 70.
- Endries M J y Adler G H. (2005) Spacing patterns of a tropical forest rodent, the spiny rat (*Proechimys semispinosus*), in Panama. *Journal of Zoology of London* 265, 147-155.
- Falk K, Benvenuti S, Dall'Antonia L, Gilchrist G y Kampp K. (2002) Foraging behaviour of thick-billed murres breeding in different sectors of the North Water polynya: an inter-colony comparison. *MARINE ECOLOGY PROGRESS SERIES* 231, 293-302.
- Fernández N. (2005) Spatial patterns in European rabbit abundance after a population collapse. *Landscape Ecology* 20, 897-911.
- Franklin W. (1983) Contrasting socioecologies of South America's wild camelids: the vicuña and the guanaco. En: M.A. Mares and N.N. Genoways (Eds), *Mammalian biology in South America. Spec Publ Ser vol 6*. Laboratory of Ecology, University of Pittsburgh, pp. 573-629.
- Fretwell S D y Lucas H L. (1970) On territorial Behavior and other factors influencing habitat distribution in birds. 16-36.
- CENPAT, CONICET, Argentina. (1982) Recuentos de línea de marcha de ñandú y mara. Contribución 65. Garrido J y Kovacs Z.
- Garthe S. (1997) Influence of hydrography, fishing activity, and colony location on summer seabird distribution in the south-eastern North Sea. *ICES Journal of Marine Science* 54, 566-577.
- Gonnet J M y Ojeda R A. (1998) Habitat use by small mammals in the arid andean foothills of the monte desert of Mendoza, Argentina. *Journal of Arid Environments* 38, 349-357.
- Greene C M y Stamps J A. (2001) Habitat selection at low population densities. *Ecology* 82(8), 2091-2100.
- Greig-Smith P. (1952) The use of random and contiguous quadrats in the study of the structure of plant communities. *Annals of Botany* 16, 293-316.
- Halama K J y Dueser R D. (1994) Of mice and habitats: tests for density-dependent habitat selection. *Oikos* 69(1), 107-114.

- Haller M, Monti A J A y Meister C M. (2001) Península Valdés. Hoja Geológica 4363-1. Programa Nacional de cartas Geológicas de la República Argentina, 24 pp.
- Hanski I y Gaggiotti O E. (2004) Ecology, genetics, and evolution of metapopulations. Elsevier.
- Hayes L D, Chesh A S y Ebensperger L A. (2007) Ecological predictors of range areas and use of burrow systems in the diurnal rodent, *Octodon degus*. *Ethology* 113, 155-165.
- Holland E P, Aegerter J N y Dytham C. (2008) Comparing resource representations and choosing scale in heterogeneous landscapes. *Landscape Ecology* 24(2), 213-227.
- Hooge P N y Eichenlaub B. (1999) Animal Movement extension to ArcView, version 2.04, beta. Anchorage, AK: Alaska Science Center, Biological Science Office, U.S. Geological Survey.
- Hoogland J L. (1981) The evolution of coloniality in white-tailed and black-tailed prairie dogs (Sciuridae: *CynomysLeucurus* and *C. Ludovicianus*). *Ecology* 62(1), 252-272.
- Hughes J y Ward D. (1993) Predation risk and distance to cover affect foraging behaviour in Namib Desert gerbils. *Animal Behaviour* 46, 1243-1245.
- Imbeau L, Drapeau P y Mönkkönen M. (2003) Are forest birds categorised as “edge species” strictly associated with edges? *Ecography* 26, 514–520.
- Ives A R y Klopper E D. (1997) Spatial Variation in Abundance Created by Stochastic Temporal Variation. *Ecology* 78, 1907-1913.
- Jensen J R. (2007) Remote sensing of the environment: an earth resource perspective. 2nd Ed., Upper Saddle River, NJ: Prentice Hall, 592 pages. 2^a ed.
- Jiao Y, Chen Y, Schneider D y Wroblewski J. (2004) A simulation study of impacts of error structure on modeling stock–recruitment data using generalized linear models. *Canadian Journal of Fisheries and Aquatic Sciences* 61, 122–133.
- Johnson C J y Seip D R. (2008) Relationship between resource selection, distribution, and abundance: a test with implications to theory and conservation. *Popul Ecol* 50, 145–157.
- Johnson M D. (2007) Measuring habitat quality: a review. *The Condor* 109, 489-504.
- Kacelnik A, Krebs J R y Bernstein C. (1992) The ideal free predator-prey distribution populations. *TREE* 7(2), 50-55.
- Keitt T H, Bivand R, Pebesma E y Rowlingson B. (2010) rgdal: Bindings for the Geospatial Data Abstraction Library. R package version 0.6-29. <http://CRAN.R-project.org/package=rgdal>.
- Keitt T H, Bjørnstad O N, Dixon P M y Citron-Pousty S. (2002) Accounting for spatial pattern when modeling organism-environment interactions. *Ecography* 5(January), 616-625.
- Kennedy M y Gray R D. (1993) Can ecological theory predict the distribution of foraging animals? A critical analysis of experiments on the ideal free distribution. *Oikos* 68(1), 158-166.
- Kraus C, Künkele J y Trillmich F. (2003) Spacing behaviour and its implications for the mating system of a precocial small mammal: an almost asocial cavy *Cavia magna*? *Animal Behaviour* 66, 225–238.
- Kufner M B y Chambouleyron M. (1991) Actividad espacial de *Dolichotis patagonum* en relación a la estructura de la vegetación en el Monte Argentino. *Studies on Neotropical Fauna and Environment* 26(4), 249-255.
- Kufner M B y Pelliza Sbriller A. (1987) Composición botánica de la dieta del mara (*Dolichotis patagonum*) y del ganado bovino en el monte mendocino. *Revista Argentina de Producción Animal* 7, 255-264.
- Lafever D H, Lafever K E, Catlin D H y Rosenberg D K. (2008) Diurnal time budget of burrowing owls in a resident population during the non-breeding season. *The Southwestern Naturalist* 53(1), 29-33.

- Legendre P y Fortin M J. (1989) Spatial pattern and ecological analysis. *Vegetation* 80, 107–138.
- León R J C, Bran D, Collantes M, Paruelo J M y Soriano A. (1998) Grandes unidades de vegetación de la Patagonia extra andina. Pp. 125-144. En: *Ecosistemas patagónicos* (Oesterheld, M.; M.R. Aguiar y J.M. Paruelo, eds.). *Ecología Austral* 8(2), 75-308.
- Levin S A. (1992) The problem of pattern and scale in ecology. *Ecology* 73, 1943-1967.
- Maher C R y Robert Burger J. (2011) Intraspecific variation in space use, group size, and mating systems of caviomorph rodents. *Journal of Mammalogy* 92(1), 54-64.
- Maier J A K, Ver Hoef J M, McGuire A D, Bowyer R T, Saperstein L y Maier H A. (2005) Distribution and density of moose in relation to landscape characteristics: effects of scale. *Canadian Journal of Forestry Research* 35, 2233-2243.
- Malizia A I. (1998) Population dynamics of the fossorial rodent *Ctenomys talarum* (Rodentia: Octodontidae). *Journal of Zoology of London* 244, 545-551.
- McGarigal K, Cushman S, Neel M y Ene E. (2002) FRAGSTATS: Spatial Pattern Analysis Program for Categorical Maps. Computer software program produced by the authors at the University of Massachusetts, Amherst. Available at the following web site: <http://www.umass.edu/landeco/research/fragstats/fragstats.html>.
- Messier F, Virgl J A y Marinelli L. (1990) Density-dependent habitat selection in muskrats: a test of the ideal free distribution model. *Oecologia* 84, 380-385.
- Mitchell M S y Powell R A. (2004) A mechanistic home range model for optimal use of spatially distributed resources. *Ecological Modelling* 177(1-2), 209-232.
- Molofsky J, Bever J D, Antonovics J y Newman T J. (2002) Negative Frequency Dependence and the Importance of Spatial Scale. *Ecology* 83(1), 21-27.
- Moody A L, Houston A I y McNamara J M. (1996) Ideal free distributions under predation risk. *Behav Ecol Sociobiol* 38, 131-143.
- Morris D W. (1987) Tests of density-dependent habitat selection in a patchy environment. *Ecological Monographs* 57(4), 269-281.
- Morris D W. (2003) Toward an ecological synthesis: a case for habitat selection. *Oecologia* 136, 1-13.
- Morris D W, Lundberg P y Ripa J. (2001) Hamilton's rule confronts ideal free habitat selection. *Proc R. Soc. Lond.* 268, 921-924.
- Nabte M J. (2010) Desarrollo de criterios ecológicos para la conservación de mamíferos terrestres en Península Valdés. Tesis Doctoral, Universidad Nacional de Mar del Plata.
- Ojeda R y Pardiñas U. (2008) *Dolichotis patagonum*. In: IUCN 2009. IUCN Red List of Threatened Species. Version 2009.1. <<http://www.iucnredlist.org/>>. Downloaded on 21 August 2009.
- Oksanen J, Blanchet F G, Kindt R, Legendre P, O'Hara R B, Simpson G L, Solymos P, Stevens M H H y Wagner H. (2010) vegan: Community Ecology Package. R package version 1.17-3. <http://CRAN.R-project.org/package=vegan>.
- Orians G H y Wittenberger J F. (1991) Spatial and temporal scales in habitat selection. *The American Naturalist* 137, 529-549.
- Oro D. (2008) Living in a ghetto within a local population: an empirical example of an Ideal Despotic Distribution. *Ecology* 89(3), 838-846.
- Overmars K P, de Koning G H J y Veldkamp A. (2003) Spatial autocorrelation in multi-scale land use models. *Ecological Modelling* 164, 257-270.
- Paradis E, Claude J y Strimmer K. (2004) APE: analyses of phylogenetics and evolution in R language. *Bioinformatics* 20, 289-290.
- Parker G A y Sutherland W J. (1985) Ideal free distributions when individuals differ in competitive ability: phenotype-limited ideal free models. *Animal behaviour* 34, 1222-1242.

- Paruelo J, Beltrán A, Jobbágy E, Sala O E y Golluscio R A. (1998) The climate of Patagonia: general patterns and controls on biotic processes. *Ecología Austral* 8, 85-101.
- Pebesma E J. (2004) Multivariable geostatistics in S: the gstat package. *Computers & Geosciences* 30, 683-691.
- Perry G L W, Miller B P y Enright N J. (2006) A comparison of methods for the statistical analysis of spatial point patterns in plant ecology. *Plant Ecology* 187(1), 59-82.
- Perry J N, Liebhold A M, Rosenberg M S, Dungan J, Miriti M, Jakomulska A y Citron-Pousty S. (2002) Illustrations and guidelines for selecting statistical methods for quantifying spatial pattern in ecological data. *Ecography* 25(January), 578-600.
- Pinheiro J, Bates D, DebRoy S, Sarkar D y R Core team. (2009) nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-96.
- Prohaska F. (1976) The climate of Argentina, Paraguay and Uruguay. En: W. Schwertfeger (Ed), *Climates of Central and South America*, pp. 13-112. Vol. 12. Elsevier, Amsterdam.
- Quirici V, Castro R A, Ortiz-Tolhuysen L, Chesh A S, Robert Burger J, Miranda E, Cortés A, Hayes L D y Ebensperger L A. (2010) Seasonal variation in the range areas of the diurnal rodent *Octodon degus*. *Journal of Mammalogy* 91(2), 458-466.
- R Development Core Team. (2010) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- Raedeke K. (1979) Population dynamics and socioecology of the guanaco (*Lama guanicoe*) of Magallanes, Chile. Doctoral, Univ. of Washington, Seattle, USA.
- Rayor L S. (1985) Effects of habitat quality on growth, age of first reproduction, and dispersal in Gunnison's prairie dogs (*Cynomys gunnisoni*). *Canadian Journal of Zoology* 63(12), 2835-2840.
- Reeves M C. (2008) Estado de conservación del mara (*Dolichotis patagonum*) en el Área Natural Protegida Península Valdés. Tesis de Licenciatura, Universidad Nacional de la Patagonia San Juan Bosco, Puerto Madryn.
- Rhodes J R, McAlpine C A, Zuur A F, Smith G M y Ieno E N. (2009) GLMM applied on the spatial distribution of koalas in a fragmented landscape. En: A.F. Zuur, E.N. Ieno, N.J. Walker, A.A. Saveliev and G.M. Smith (Eds), *Mixed effects models and extensions in ecology with R*. Springer, Nueva York.
- Ringuelet R A. (1961) Rasgos fundamentales de la zoogeografía de la Argentina. *Physis* 22, 151-170.
- Ripley B D. (1976) The second-order analysis of stationary point processes. *Journal of Applied Probability* 13, 255-266.
- Ripley B D. (1988) *Statistical inference for spatial processes*. Cambridge University Press.
- Rodríguez D. (2009) Modeling habitat use of the threatened and endemic mara (*Dolichotis patagonum*, Rodentia, Caviidae) in agricultural landscapes of Monte Desert. *Journal of Arid Environments* 73, 444-448.
- Rodríguez D y Dacar M A. (2008) Composición de la dieta de la mara (*Dolichotis patagonum*) en el sudeste del monte pampeano (La Pampa, Argentina). *Mastozoología Neotropical* 15(2), 215-220.
- Rodríguez M D. (2003) Caracterización de las áreas de uso de la mara *Dolichotis patagonum* en el sudeste de La Pampa. Tesis de Licenciatura en Ciencias Biológicas, Universidad Nacional de La Pampa.
- Rosenzweig M L. (1991) Habitat selection and population interactions: the search for mechanism. *The American Naturalist* 137, S5-S28.
- Rosenzweig M L. (1981) A theory of habitat selection. *Ecology* 62(2), 327-335.

- Roshier D a, Doerr V a J y Doerr E D. (2008) Animal movement in dynamic landscapes: interaction between behavioural strategies and resource distributions. *Oecologia* 156(2), 465-477.
- Rostagno C M. (1981) Propiedades, génesis y clasificación de dos suelos de Península Valdés, Provincia del Chubut. Actas IX RACS y "Reconocimiento de los Suelos de Península Valdés" contribución 44 – CENPAT.
- Rowe D L y Honeycutt R L. (2002) Phylogenetic relationships, ecological correlates, and molecular evolution within the Caviioidea (Mammalia, Rodentia). *Molecular Biology and Evolution* 19, 263-277.
- Sachser N. (1985) Different forms of social organization at high and low population densities in guinea pigs. 253-271.
- Schaefer J A y Messier F. (1995) Habitat selection as a hierarchy: the spatial scales of winter foraging by muskoxen. *Ecography* 18, 333-344.
- Schiffers K, Schurr F M, Tielbörger K, Urbach C, Moloney K y Jeltsch F. (2008) Dealing with virtual aggregation - a new index for analysing heterogeneous point patterns. *Ecography* 31, 545-555.
- Schooley R L y Wiens J A. (2001) Dispersion of kangaroo rat mounds at multiple scales in New Mexico, USA. *Landscape Ecology* 16, 267-277.
- Schurr F M, Bossdorf O, Milton S J y Schumacher J. (2004) Spatial pattern formation in semi-arid shrubland : a priori predicted versus observed pattern characteristics. *Plant Ecology* 173, 271-282.
- Sinclair A R E. (1992) Do large mammals disperse like small mammals? En: N.C. Stenseth and W.Z. Lidicker (Eds), *Animal dispersal. Small mammals as a model*, pp. 229-242. Chapman & Hall, London.
- Skellam J G. (1951) Random dispersal in theoretical populations. *Biometrika* 38(1-2), 196-218.
- Sombra M S y Mangione A M. (2005) Obsessed with grasses? The case of mara *Dolichotis patagonum* (Caviidae: Rodentia). *Revista Chilena de Historia Natural* 78, 401-408.
- Stamps J A, Krishnan V V y Reid M L. (2005) Search costs and habitat selection by dispersers. *Ecology* 86(2), 510-518.
- Stapp P. (1997) Habitat selection by an insectivorous rodent: patterns and mechanisms across multiple scales. *Journal of Mammalogy* 78(4), 1128-1143.
- Stephens P A y Sutherland W J. (1999) Consequences of the Allee effect for behaviour, ecology and conservation. *Trends in Ecology & Evolution* 14, 401-405.
- Stoyan D y Penttinen A. (2000) Recent applications of point process methods in forestry statistics. *Statistical Science* 15(1), 61-78.
- Tabeni S, Mastrantonio L y Ojeda R A. (2007) Linking small desert mammal distribution to habitat structure in a protected and grazed landscape of the Monte, Argentina. *Acta Oecologica* 31, 259-269.
- Tabeni S y Ojeda R A. (2003) Assessing mammal responses to perturbations in temperate aridlands of Argentina. *Journal of Arid Environments* 55, 715-726.
- Taber A B. (1987) The behavioural ecology of the mara *Dolichotis patagonum*. Ph.D. dissertation., University of Oxford.
- Taber A B y MacDonald D W. (1992a) Communal breeding in the mara, *Dolichotis patagonum* (Rodentia: Caviomorpha). *Journal of Zoology of London* 227(439-452).
- Taber A B y MacDonald D W. (1992b) Spatial organization and monogamy in the mara *Dolichotis patagonum*. *Journal of Zoology of London* 227, 417-438.
- Taraborelli P, Corbalan V y Giannoni S. (2003) Locomotion and escape modes in rodents of the Monte Desert (Argentina). *Ethology* 109(6), 475-485.
- Tregenza T, Parker G A y Thompson D J. (1996) Interference and the ideal free distribution: models and tests. *Behavioral Ecology* 7(4), 379-386.

- Valone T J. (2007) From eavesdropping on performance to copying the behavior of others: a review of public information use. *Behavioural Ecology and Sociobiology* 62, 1-14.
- van Teeffelen A J A y Ovaskainen O. (2007) Can the cause of aggregation be inferred from species distributions? *Oikos* 116, 4-16.
- Venables W N y Ripley B D. (2002) *Modern Applied Statistics with S*. Fourth Edition ed. Springer, Nueva York.
- Wahlström K y Kjellander P. (1995) Ideal free distribution and natal dispersal in female roe deer. *Oecologia* 103, 302-308.
- Watt A S. (1947) Pattern and process in the plant community. *Journal of Ecology* 35, 1-22.
- Wilson T L, Rayburn A P y Edwards T C, Jr. (2012) Spatial ecology of refuge selection by an herbivore under risk of predation. *Ecosphere* 3(1), <http://dx.doi.org/10.1890/ES11-00247.1>.
- Zuur A F, Ieno E N, Walker N J, Saveliev A A y Smith G M. (2009) *Mixed Effects Models and Extensions in Ecology with R*, 554 pp. *Statistics for Biology and Health*, edited by M. Gail, K. Krickeberg, J.M. Samet, A. Tsiatis and W. Wong. Springer, New York.