

BIBLIOGRAFÍA

Andrade F., Cirilo A., Uhart S., Otegui M. (1996). Introducción, Características Del Cultivo de Maiz. En: Ecofisiología del Cultivo de Maíz. (La Barrosa, ed.) Balcarce, Buenos Aires, Argentina.

Aulinger I. E. (2002). Combination of in Vitro androgenesis and biolistic transformation: an approach for breeding transgenic maize (*Zea Mays L.*) lines. Dissertación para optar por el grado de doctor en Ciencias Naturales Swiss Federal Institute Of Technology Zurich.

Ayres Guidetti Zagatto Paterniani M. E. (2001). Use of heterosis in Maize Breeding: History, Methods and Perspectives-A Review. *Crop Breeding and Applied Biotechnology* v.1 n.2, p159-178.

Baenziger P. S., Russell W.K., Graef G.L. and Campbell B.T. (2006) Improving lives: 50 years of Crop Breeding, Genetics and Cytology (C-1). *Crop Sci.* 46:2230-2244.

Barclay I. R. (1975) High frequencies of haploid production in wheat (*Triticum aestivum*) by chromosome elimination. *Nature* 256: 410-411.

Barnabás B., Obert B., Kovacs G. (1999). Colchicine, an efficient genome-doubling agent for maize (*Zea mays L.*) microspores cultured in anthero. *Plant Cell Reports* 18:858-862.

Belicuas P.R. (2004). Obtención, identificación e caracterización de haplóides androgenéticos em milho. Universidad Federal de Lavras. Magister em Agronomía, Minas Gerais, Brasil. pg 52.

Berger, F. (1999) Endosperm development. Curr. Op. Plant Biol. (2):28-32.

Bordes J., Charmet G., Dumas de Vaulx R., Pollacsek M., Beckert M., Gallais A. (2006). Doubled haploid versus S1 family recurrent selection for testcross performance in a maize population. Theor. Appl Genet 112:1063-1072.

Bouvier L., Fillon F. R., Lespinasse Y. (1994). Oryzalin as an efficient agent for Chromosome Doubling of Haploid Apple Shoots in vitro Plant Breeding 113: 343-346.

Brazauskas G., Padakinskienë I., Ruzgas V. (2005). Improved approaches in wheat x maize crossing for wheat doubled haploid production. Biologija Nr. 4. P. 15-18.

Castillo A. M., Vallés M. P. & Cistué L.. 2000. Comparison of anther and isolated microspore cultures in barley. Effects of culture density and regeneration medium. Euphytica. 113:1-8.

Coe E.H.Jr y Sarkar K.R. (1964). The detection of haploids in maize. Journal of Heredity (55): 231-233.

Coe E. H. Jr., Neuffer M. G., Hoisington D. A. (1988). The Genetics of Corn. En: Corn and corn Improvement 3ra Edición. (Sprague C. F., Dudley J. W. eds) pg 81-258. A.S.A., C.S.S.A., S.S.S.A, Madison Winsconsin, USA, 986 pg.

Comeau, A., Nadeau P., Plourde A., Simard R., Maës O., Kelly S., Harper L., Lettre J., Landry B., and C. A. St. Pierre. (1992). Media for the in ovule culture of protoembryos of wheat and wheat-derived interespecific hybrids or haploids. Plant Science 81:117-125.

Costa Rodriguez M.. D'Ayala Valva F., Madureira Brasil E., Chaves Lázaro J. (2001). Comparison among inbreeding systems in maize. Crop Breed. And Appl. Biotech. (1) 2: 105-114.

Chalyk S.T. (1994). Properties of maternal haploid maize plants and potencial application to maize breeding. Euphytica 79:13-18.

Chalyk S., Baumann A., Daniel G., Eder J. (2003). Aneuploidy as a possible cause of haploid-induction in maize. En: Maize Genetics Cooperation Newsletter, USA, (Maize Genetics Cooperation Newsletter, ed), Vol 77.

Chalyk S.T., Bylich V.G. and Chebotar O.D. (1994). Transgressive segregation in the progeny of a cross between two inducers of maize maternal haploids. Maize genetics Database. <http://www.maizegdb.org/mnl/68/69chalyk.html>

Chase S., (1969) Monoploids and monoploid-derivatives of maize (*Zea mays* L.) The Botanical Review (35):119-120.

Deimling S., Röber F., Geiger H.H. (1997). Methodik und Genetik der in-vivo-Haploidinduktion bei Mais. Vortr Pflanzenzüchtung 353:225-226

De Laat A.M.M., Gohde W., Vogelzang M.J.D.C. (1987). Determination of Ploidy of Single Plants and Plant Populations by Flow Cytometry. Plant Breeding 99:303-307.

Eder J., Chalyk S. (2002). In vivo haploid induction in maize. Theor. Appl. Genet. (104):703-708.

Eder J., Chalyk S. (2003). In-vivo induktion von Haploiden-praktische Anwendung in der Linientwicklung und rekurrenten Selektion bei Mais. Tagung 2003 der Vereinigung der Pflanzenzüchter und Saatgutkaufleute Österreichs.(54):1-3.

Eyherabide G.H., Lorea R.D., Delucchi C., López C.G., Ferrer M.E., Presello D.A., Defacio R. Evaluación del potencial de variedades nativas de maíz como recurso para el mejoramiento de un cruzamiento entre líneas duras coloradas. En: Actas del VIII Congreso Nacional de Maíz 16-18 Noviembre 2005., Pergamino, Argentina, 463 pg.

East E.M. y Hayes H.K. (1911). Inheritance in maize. Connecticut Agric. Exp. Stn. Bull. 167

Friedt, W., Foroughi-Wehr (1981). Anther culture of barley. Plant regeneration and agronomic performance of diploid progenies. En: Actas IV International Barley Genetics Symposium, (M Asher, ed). Edinburgh Univ. Press, Edinburgh. Edinburgo, Escocia, P. 690-698

Fukuyama T., Hosoya H. (1983). Genetic control and mechanism of chromosome elimination in the hybrids between *Hordeum bulbosum* (4X) and *H. vulgare* (4X). Jpn. J. Genet 58: 241-250.

Gear J.R.E. (2006). El Cultivo del Maíz en la Argentina. En: Maíz y Nutrición, Informe sobre los usos y las propiedades nutricionales del maíz para la alimentación humana y animal. Recopilación de Internacional Life Sciences Institute (ILSI), Argentina. Serie de Informes especiales. www.maizar.org.ar, 80 pg.

Germand D., Rutten T., Varshney A., Rubtsova M., Prodanovic S., Brüs C., Kumlehn J., Matzk F., Houben A. (2005) Uniparental chromosome elimination at mitosis and interphase in wheat and pearl millet crosses involves micronucleus formation, progressive heterochromatinization, and DNA fragmentation. The Plant Cell, Vol. 17, 2431–2438

Goodman M. M., Brown W. L. (1988). The Races of Corn. En: Corn and corn Improvement 3ra Edición. (Sprague C. F., Dudley J. W. eds) pg 81-258. A.S.A., C.S.S.A., S.S.S.A, Madison Winsconsin, USA, 986 pg.

Hallauer A.R., Russell W.A., Lamkey K.R. (1988). Corn Breeding. En: Corn and corn Improvement 3ra Edición. (Sprague C. F., Dudley J. W., eds) pg 463-564. A.S.A., C.S.S.A., S.S.S.A, Madison Winsconsin, USA, 986 pg.

Hansen N.J.P, Andersen S.B. (1998). Efficient production of doubled haploid wheat plants by in vitro treatment of microspores with trifluralin or APM. Plant Breeding 117:401-405.

Ho I., Wan Y., Widholm J.M. and Rayburn A. L. (1990). The use of stomatal chloroplast number for rapid determination of ploidy level in maize. Plant Breeding 105: 203-210.

Hu T., Kasha K.J. (1997). Improvement of isolated microspore culture of wheat (*Triticum aestivum* L) through ovary co-culture. Plant Cell Reports 16: 520-525.

Huang B.Q., Sheridan W.F. (1996) Embryo sac development in the maize *indeterminate gametophyte1* mutant: abnormal nuclear behavior and defective microtubule organization. The Plant Cell (8): 1391-1407.

Huang B., Sunderland, N. (1982). Temperature-stress Pretreatment in barley anther culture. Ann Bot. 49: 77-88.

InfoStat (2008). *InfoStat versión 2008*. Grupo InfoStat, FCA, Universidad Nacional de Córdoba, Argentina.

Jugenheimer R.W. (1976) Hybrid Corn Breeding Systems. En:Corn improvement, seed production and uses. pg 93-110. John Wiley and Sons, Wiley interscience publication, USA.

Kasha K., Kao. K. N. (1970). High frequency haploid production in barley. Nature 225:874-876.

Kasha K. J., Simion E., Oro R., Yao Q.A., Hu T. C. & Carlson A. R. (2001). An improved in vitro technique for isolated microspore culture of barley. Euphytica. 120: 379-385.

Kato A. (1997). Induced single fertilization in maize. Sex. Plant Reprod. (10):96-100.

Kato A. (2002) Chromosome doubling of haploid maize seedlings using nitrous oxide gas at the flower primordial stage. Plant Breeding 121: 370-377.

Kato A. and Birchner J. A. 2006 Induction of tetraploid derivatives of Maize Inbred lines by Nitrous Oxide Gas treatment. Journal of Heredity 97:39-44.

Kermicle J.L. (1969). Androgenesis conditioned by a mutation in maize. Science (166): 1422-1424.

Kermicle J.L. (1971). Pleiotropic effects on seed development of the indeterminate gametophyte gene in maize. Amer. J. Bot. 58(1):1-7

Kraptchev B., Kruleva M., Dimitrov B., Markova M. (2001). Investigations of sources for haploidy in maize (*Zea mays* L). *Maydica* (46):215-219.

Kyndiger B. and Hamann S. (1993). Generation of haploids in maize: a modification of the indeterminate gametophyte (ig) system. *Crop Sci.* (33): 344-346

Lashermes P., Beckert M. (1988). Genetic control of maternal haploidy in maize (*Zea mays* L.) and selection of haploid inducing lines. *Theor Appl Genet* (76):405-410

Lashermes P., Gaillard A. and Beckert M. (1988). Gynogenetic haploid plants analysis for agronomic and enzymatic markers in maize (*Zea mays* L.). *Theor Appl Genet* 76:570-572.

Laurie D.A., Bennett M.D. (1988). The production of haploid wheat plants from wheat x maize crosses. *Theor. Appl. Genet.* 76:393-397.

Longin C.F.H., Utz H.F., Reif J.C., Schipprack W., Melchinger A.E. (2006) Hybrid maize breeding with doubled haploids: I. One-stage versus two-stage selection for testcross performance. *Theor. Appl Genet* (112):903-912.

Martins-Lopes P.F., Guedes-Pinto H., Pinto-Carnide O., Snape J. (2001) The effect of spikelet position on the success frequencies of wheat haploid production using the maize cross system. *Euphytica* 121:265-271.

Orton T.J. (1980). Haploid barley regenerated from callus cultures of *Hordeum vulgare* X *H. jubatum*. The journal of Heredity 71: 280-282.

Polci P., Conti V., Aldao Humble G., Miranda R., Echenique V. (2006) Obtención de plantas haploides de cultivares argentinos de trigo pan (*Triticum aestivum*) por cultivo de anteras y cruzamientos con maíz. RIA 34 (3): 151-176

Polci P., Conti V., Miranda R., (2004). Obtención de plantas doblehaploides. En: Métodos para acelerar programas de mejoramiento e identificación varietal. IV-Capítulo 1. Biotecnología y Mejoramiento Vegetal. (Viviana Echenique, Clara Rubinstein, Luis Mroginski, eds). Ediciones INTA, 2004, Buenos Aires, Argentina, 448 pg.

Röber F.K., Gordillo G.A., Geiger H.H. (2005) In vivo haploid induction in maize- performance of new inducers and significance of doubled haploid lines in hybrid breeding. Maydica (50): 275-283.

Ryan A.B., Castillo A. M., Vallés M.P., Sanz J.M. (1999). Dried doubled-haploid embryos obtained from microspore culture of barley cv. Igri. Plant Cell Reports 18: 924-928.

San Noeum. (1976) Haploides d'*Hordeum vulgare* L. Par culture in vitro d'ovaires non fecondés. En: Barley, Barley breeding (D.C. Rasmusson, ed). American Society of Agronomy, Madison WI. p. 231-268.

Sarkar K. R., Coe E. H. (1966). A genetic Analysis of the origin of maternal haploids in maize. Genetics 54: 453-464.

Savaskan C., Szarejko I., Toker M.C. (1999). Callus production and plant regeneration from anther culture of some Turkish barley cultivars. Turkish Journal of Botany 23:359-365.

Séguin-Swartz G., Kott L., Kasha K.J. (1984) Development of haploid cell lines from immature barley, *Hordeum vulgare*, embryos. Plant Cell Reports 3:95-97.

Sendra G.H., Salerno J.C., Kandus M.V., Weber C., Arizaga R., Rabal H., Trivi M. (2005). Análisis de distintos tipos de maíz (*Zea mays* L.) mediante un método óptico (biospeckle laser) para diferenciar su calidad industrial. En: Actas del VIII Congreso Nacional de Maíz 16-18 Noviembre 2005., Pergamino, Argentina, 463 pg.

Shatskaya O.A., Zabirova E.R., Shcherbak V.S., Chumak M.V. 1994. Mass induction of maternal haploids in corn. Maize Gen. Coop. Newsletter 68:51.

Shull G.H. (1908). The composition of a field of maize. Am. Breeder's Assoc. Rep. 4:296-301.

Shull G.H. (1909). A pure-line method in corn breeding. Am. Breeder's Assoc. Rep. 5:51-59.

Shull G.H. (1910). Hybridization methods in corn breeding. Am. Breeder's Mag. 1:98-107.

Strickberger M.W. (1998). Cambio de las frecuencias génicas. En: Genética 3ra edición. (Omega s.a., ed) Plató 26-08006 Barcelona. ISBN:84-282-0829-8.

Valla J.J. (2000). Flor. Megagametogénesis. En: Botánica. Morfología de las plantas superiores (Hemisferio Sur, ed), Buenos Aires, Argentina, 332 pg.

Wan Y., Widholm J.M. (1995). Effect of chromosome-doubling agents on somaclonal variation in the progeny of doubled haploids of maize. Plant Breeding 114: 253-255

Wang C. C., Kuang B. J. (1981). Induction of haploid plants from the female gametophyte of *Hordeum vulgare* L. In Barley Breeding. P.231-268. 56 in: Barley. D.C. Rasmusson, ed. American Society of Agronomy, Madison WI.

Zhou H., Konzak C.F. (1989). Improvement of anther culture methods for haploid production in wheat. Crop Science 29:817-821